

CITY OF LAKE FOREST PARK SPECIAL TREE BOARD MEETING

Wednesday, November 29, 2023 at 6:00 PM

Meeting Location: In Person and Virtual / Zoom 17425 Ballinger Way NE Lake Forest Park, WA 98155

INSTRUCTIONS FOR PARTICIPATING IN THIS MEETING VIRTUALLY:

Join Zoom Webinar: https://us06web.zoom.us/j/88299231947 Call into Webinar: 253-215-8782 | Webinar ID: 882 9923 1947

The Tree Board is providing opportunities for public comment by joining the meeting webinar (via computer or phone) or in person to provide oral public comment.

HOW TO PARTICIPATE WITH ORAL COMMENTS:

If you are attending in person, there is a sign-in sheet located near the entrance to the room. Fill out the form and the presiding officer will call your name at the appropriate time. Oral comments are limited to 3:00 minutes per speaker.

If you are attending the meeting via Zoom, in order to address the Tree Board during the Public Comment section of the agenda, please use the "raise hand" feature at the bottom of the screen. Oral comments are limited to 3:00 minutes per speaker. Individuals wishing to speak to agenda items will be called to speak first in the order they have signed up. The meeting host will call your name and allow you to speak. Please state your name and whether you are a resident of Lake Forest Park. The meeting is being recorded.

For up-to-date information on agendas, please visit the City's website at www.cityoflfp.gov

AGENDA

- 1. CALL TO ORDER: 6:00 PM
- 2. SHORT REFLECTION
- 3. INTRODUCTIONS
- 4. ADOPTION OF AGENDA
- 5. APPROVE MINUTES
 - A. July 5, 2023 Tree Board Meeting Minutes
 - B. October 4, 2023 Tree Board Meeting Minutes
- 6. CITIZEN COMMENTS

This portion of the agenda is set aside for the public to address the Tree Board on agenda items. **Comments are limited to a three (3) minute time limit.**

7. COMMUNICATION

8. OLD BUSINESS

- A. Website and Social Media Outreach
- B. Proposed Flyer for City Newsletter

9. **NEW BUSINESS**

A. Draft Urban Forest Ecosystem Sevices and Values Report from DCG/Watershed

10. REPORTS AND ANNOUNCEMENTS

A. November Arborist Report

11. AGENDA FOR NEXT MEETING

12. ADJOURN

Any person requiring a disability accommodation should contact city hall at 206-368-5440 by 4:00 p.m. on the day of the meeting for more information.

City of Lake Forest Park – Tree Board Meeting Normal Meeting Minutes: July 5, 2023; 7:00-9:00pm Hybrid Meeting Held in the Forest Room at City Hall and Virtually via Zoom

Tree Board Members present: Chair Richard Olmstead, Boardmembers Mark Phillips, Marty Byrne, and Doug Sprugel

Staff and others present: Riley Bushnell, Assistant Planner; Councilmember Larry Goldman

Members of the Public present: Scott Morrison, Nancy Hertzog, Julie Turnell, Randi Sibonga

Tree Board Members absent: Board Member Sandra LeVar

Call to order: 7:02 PM

Short Reflections: Board Member Phillips gave a short reflection.

Introductions: No need for introductions

Approval of Meeting Agenda: Board Member Phillips moved to approve the agenda. Board Member Sprugel seconded the motion, and the motion to approve the agenda carried.

Approval of Minutes: Board Member Phillips moved to approve the minutes. Board Member Sprugel seconded the motion, and the motion to approve the minutes carried.

Public Comment: None.

Communication: The Board discussed possible improvements to the tree removal permit report and the retirement of Planning Director Stephen Bennett.

Old Business:

Priority areas for plantings

The Board went over a map of possible priority areas for possible planting of new trees related to the possible tree removals in a right-of-way corridor.

Right-of-way corridor permit regulations

The Board discussed the proposed amendments to the Tree Code.

The Board discussed the possible design of Lakefront Park.

Education Outreach

Board Member Byrne gave a brief update on the education outreach and the possible demonstration regarding ivy removal in the future. The Board also discussed possible improvements for the city website and attendance at future events.

The Board continued to discuss the possible tree corridor in the right of way along SR 522.

New Business:

Reports and Announcements:

Tree Permit Report

The Assistant Planner passed around copies of Arborist Swanson's monthly report for May tree permits.

Agenda for Next Meeting:

A new agenda order was discussed for the next meeting.

Adjournment: Board Member Sprugel moved to adjourn. Board Member Phillips seconded; the meeting adjourned at 8:20 p.m.

1 2 3 4	City of Lake Forest Park – Tree Board Meeting Normal Meeting Minutes: October 4, 2023; 7:00-9:00pm Hybrid Meeting Held in the Forest Room at City Hall and Virtually via Zoom
5 6 7	Tree Board Members present: Chair Richard Olmstead, Board Members Mark Phillips, Doug Sprugel, Marty Byrne, and Sandra LeVar (via Zoom)
8 9	Staff and others present: Matt McLean, City Clerk; Councilmember Larry Goldman
10 11	Members of the Public present: John Drew, Sara Phillips
12 13	Tree Board Members absent: None.
14 15	Call to order: 7:00 PM
16 17	Introductions: The Tree Board members and City staff introduced themselves.
18	Short Reflections: Chair Olmstead gave a short reflection.
19 20 21 22	Approval of Meeting Agenda: Chair Omstead moved to approve the agenda. There was on objection, and it was considered approved.
23 24	Approval of Minutes: There were no minutes for approval.
25	Public Comment:
26 27	• Sara Phillips commented on the urban forest planting event that is occurring at Shoreline Historical Museum on December 9th.
28 29	John Drew commented on the meeting he attended from Sound Transit
30	Communication:
31 32	The Board discussed matters regarding staffing in the city.
33	Old Business:
34	Ivy Removal Demonstration
35	The Board discussed the ivy removal demonstration and the recording of a video showing how to
36	properly remove ivy from trees.
37 38	Draft Ordinance regarding amendments to the Tree Code related to the preservation of the tree
39	canopy
40	The Board discussed the current draft of the amendments to the Tree Code and the need for an
41 42	updated tree list.
43	Strategy to search for new Tree Board members
44	The Board discussed options to advertise for new Tree Board members including using the city's
45	social media.
46	
47	

1	
2	New Business:
3	Website update proposal from Board Member LeVar
4	Board Member LeVar reviewed possible improvements that can be made to the city website related
5	to the Tree Board and tree permits.
6	1
7	Arborist/Permit Information from Board Member LeVar
8	Board Member LeVar reviewed possible flyer information regarding recruiting for an arborist, tree
9	board members, and when people need to obtain a tree permit.
10	
11	Planning for the next canopy study
12	Chair Olmstead wanted to keep the Board aware that the canopy study will need to be completed in
13	the future.
13 14 15	
15	Reports and Announcements:
16	Active Tree Permit Reports
17	The Board discussed the active tree permit reports.
18	
19	Agenda for Next Meeting:
20 21	The Board discussed the need to review the tree inventory study at a future meeting.
21	
22	Adjournment: The meeting adjourned at 9:00 PM.
23	
24	
25	APPROVED:
26	
22 23 24 25 26 27 28	
28	Richard Olmstead, Chair

Tree Board - Education Outreach

MEDIA & CONTENT PROPOSAL

By: Marty Byrne & Mandee Kulaga Parker with Sandra LeVar - Lake Forest Park Tree Board Members

Media - EDUCATION	1
Content - Education	2
Audience - Education & OUTREACH	3
Understanding LFPs Municipal Code / Trees	Э
Introducing the Tree Board	3
Content for newsletters	4
APPENDIX – Understanding the Tree Code	5

Proposal for 2023-2024 outreach education for the Lake Forest Park Tree Board includes a staged approach to media content. Goals include maintaining the city canopy and increasing public satisfaction. Approach for first phase includes reorganization of website content followed by updated website content. Initial target audience includes existing residents. Priorities are:

- Community outreach for new board members needed asap
- One-time educational outreach mailer introducing LFP Tree Board
- · One-time educational outreach mailer introducing lvy Out
- One-time redesign of tree board pages on LFP website
- · On-going content updates on LFP Tree Board pages
- Regular quarterly/monthly content contributions to eNewsletter and social media
- As needed posters, flyers, and handouts for events:
 - Arbor Day
 - Green Day
 - 0 ...

MEDIA - EDUCATION

Website

- Inventory existing content on LFP website related to trees and the canopy
- Outline preferred content for Tree Permit (and FAQ) and Tree Board webpages

Page 2 of 8 DRAFT AI Section 8, ItemA.

Draft educational content for sections of Tree related webpages

eNewsletter & LFP Times

- · Educational content to be included in monthly eNewsletter
- Content in eNewsletter to be summarized for quarterly LFP Times

Social Media Infographics

Content from Website and eNewsletters to be incorporated into infographics

CONTENT - EDUCATION

Permit How to & Why (for landowners)

https://www.cityoflfp.com/163/Tree-Permits-and-Information

Include code changes

Permit Approved & Open for Comments (for residents)

Why & How to comment

The Canopy & Why it's a "Good Thing" (for landowners and developers)

... TBD ...

Introduction to Tree Services (for resident and landowners)

List of Tree Services/Contractors that have submitted/approved signed "Statement of Canopy Preservation and Enhancement Acknowledgement" form with the Business License Application

Good Trees to Plant

https://www.cityoflfp.com/239/Tree-List

Add new updated information about how to plant new trees; and include information from local PNW university websites and research. Links include:

- WSU, Oregon SU, UC Davis
- City of Seattle, City of Kirkland
- WSU Extension: Backyard Forest Stewardship in Western Washington (pub EM107E)
- King Conservation District: Backyard Forest Stewardship Homeowners Guide
- Arbor Day Foundation: ... < lots to choose from>

What are good trees to plant? What are good places to plant trees?

Page 3 of 8 DRAFT A Section 8, ItemA.

Taking Care of Your Trees

https://www.cityoflfp.com/239/Tree-List

Add new updated information about how to maintain existing trees; and include information from local PNW university websites and research.

Walk about (trees and parks)

https://www.cityoflfp.com/240/Tree-Walks-of-Lake-Forest-Park

AUDIENCE - EDUCATION & OUTREACH

Residents

Anyone who lives in Lake Forest Park - landowners or renters

Businesses

Landowners or leaseholders of retail space/land in Lake Forest Park

Landowners

Commercial or Residential landowners that may or may not live in Lake Forest Park

Developers

Businesses owners that improve land including construction, landscape, masonry, etc

Tree Services

Businesses that inspect and or maintain trees

UNDERSTANDING LFPS MUNICIPAL CODE / TREES

https://www.cityoflfp.com/fag.aspx?TID=16

Chapter 16.14 Tree Canopy Preservation and Enhancement

INTRODUCING THE TREE BOARD

- Effort owned by:
- Content for letter
- Mailing list
- Printer
- Mail Merge & Send

Welcome / Introduction to Tree Board (What can/do we do for YOU — residents, landowners, developers)

Page 4 of 8 DRAFT A Section 8, ItemA.

Welcome to Lake Forest Park. This is a fantastic community; everyone loves it here and we hope you will also. We care about our neighbors and respect the Urban Forest that surrounds us. We have safe streets and pleasant green parks that offer welcoming places to walk and meet neighbors.

Ole Hanson developed Lake Forest Park circa 1912 "with an eye on Nature." He wished to maintain a city where people enjoyed "the stately cedar, the majestic fir, the quivering cypress, and the homelike maple."

The Lake Forest Park Tree Board and the Stewardship Foundation are here to help you get better acquainted with the Lake Forest Park Urban Forest, its benefits, and the local ordinances that protect it.

Why Trees Matter (in LFP, aka Benefits of Trees, vegetation, and the Tree Board)

How our Urban Forest of healthy trees helps us:

- Enhance the beauty of one's home and fosters a pleasant place to live
- Increase the value \$\$\$\$ of one's property
- · Reduce the urban heat island effect
- Save energy cost \$\$\$\$ for homeowners
- · Promote a calming environment for the homeowner and family
- Help manage dispersal of storm water, and improve the water quality of our
- salmon rich streams
- Offer quiet green spaces fostering good psychological and physical health
- Provide habitat for wildlife and sustain biodiversity
- Sequester carbon to help slow global warmingMaintenance of Private Property Trees (How to, includes removal/replace)

Growth / Planting / Improvement of Private Property Trees (How to)

https://www.cityoflfp.com/235/Learn-More-about-Trees-Forest-Care

CONTENT FOR NEWSLETTERS

Primary objective for outreach content will be to use existing educational material, composed for a targeted audience.

- Effort Owned by:
- Schedule, Contact info for publishing
 - eNewsletter
 - Social Media
- Content/ Writer

EXAMPLES of Educational Material

- Why to submit a permit, How to submit a permit
- · Why tree removal/replacement permits are approved, How to provide public comment
- What is The Canopy, Why is it a "Good Thing"

Page 5 of 8 DRAFT AI Section 8, ItemA.

EXAMPLES of Target Audience

- Residents and Businesses
- Landowners
- Developers and Tree Services

APPENDIX – UNDERSTANDING THE TREE CODE

"Tree" means a self-supporting woody plant typically reaching at least 12 to 15 feet in height at maturity. "Significant tree" means a tree six inches or greater in diameter.

Tree Permit Types

A tree permit should be submitted by landowner (or on behalf of landowner) before removal of a tree in Lake Forest Park (public, private, residential or commercial)

https://www.cityoflfp.com/DocumentCenter/View/6247/Tree-Permit-Application?bidId=

Minor Permit

 include, but are not limited to: additions, enlargements, or alterations to existing structures, construction of retaining walls, fences, driveways, and garages, clearing and grading activity

Major Permit

- means subdivision or short subdivision; construction or demolition of single-family, multifamily, or commercial buildings; and alterations, repairs, enlargements or additions that add 1,000 square feet or more of impervious surface coverage.
- Proactive Forest Management Permit
- Utility Forest Management Permit

Tree Permit Workflow

- Submit Permit
- Review of Tree Types on permit
 - Significant Trees
 - "Significant tree" means a tree six inches or greater in diameter (DBH) or a required replacement tree of any size. Dead trees shall not be considered significant trees.
 - "Landmark tree" means a significant tree that is at least 24 inches in diameter (DBH).
 - Exceptional Trees

- Removal of viable exceptional trees, is prohibited.
- "Exceptional tree" means a viable tree, which because of its unique combination of size and species, age, location, and health is worthy of long-term retention, as determined by the city's qualified arborist.

Table 1: Exceptional Tree Species and Their Threshold Diameters					
Species	Threshold Diameter				
Bigleaf MAPLE – Acer macrophyllum	42 inches				
Douglas FIR – Pseudotsuga menziesii	42 inches				
Grand FIR – Abies grandis	33 inches				
MADRONA – Arbutus menziesii	12 inches				
Western HEMLOCK - Tsuga heterophylla	36 inches				
Western Red CEDAR - Thuja plicata	42 inches				
Western White PINE - Pinus monticola	36 inches				

- Review of Replacement Trees on permit
 - "General tree list" means a list of tree species that is maintained by the city and approved by the city's qualified
 arborist for planting as replacement trees, as well as tree species that are prohibited from being planted as
 replacement trees.
- Review of lot Canopy Coverage
 - "Canopy" means the part of the tree crown composed of leaves and small twigs or the collective branches and foliage of a group of trees 'crowns. "Canopy coverage" means the area covered by the canopy of trees on the lot.
 - A minor tree permit shall be granted if conditioned on at least one tree replacing each tree removed, to provide canopy coverage equal to or greater than the tree(s) being removed.
 - When the proposed tree removal is associated with major development activity, the trees may be removed if a
 tree replacement plan is approved that, at a minimum, brings canopy coverage to the applicable canopy
 coverage goal.

- Development Proposal Requirements
- Review of Exceptions
 - Major Permits
 - · Property that has undergone major development activity in the past five years
 - Tree Conservation Easement & Environmentally Critical Areas
- Notice of Decision: Approved, Conditioned, Denied
 - A conditional use permit is a zoning exception status.
- Appeals
- Amendments
- Expiration & Close Permit

Tree Maintenance

- Coverage
- Health
- Pruning

Enforcement

- Violations
 - In addition to tree replacement, the administrator shall require that the persons found in violation of this chapter or the conditions of a permit pay the appraised value of the trees
- Stop Work Order
 - the building official may suspend some or all of the work as appropriate
- Remedial Measures
 - The persons found in violation of the conditions of a tree removal permit or in violation of this chapter may be required to perform remedial measures as ordered by the administrator that are necessary to correct the violation

Tree Services

- LFP Business License
- A Statement of Acknowledgement

Print Media: 1/3 page double sided

LAKE FOREST PARK Washington

This is a fantastic community; and *
we know you'll love it here. We
care about our neighbors and
respect the Urban Forest that
surrounds us.



Tree Permits let home owners easily remove problem trees with the provision that a new tree be replanted



Permits help balance removals so that the neighborhood tree canopy remains balanced



GETTING TO KNOW YOUR CITY'S ARBORIST




Helping the community to protect trees for future generations.

Managing ongoing challenges including:

- Civic construction projects
- Drought, aging trees, disease
- Seasonal maintanance issues

"Ensure tree safety, value, and beauty".







Do I need to submit a Tree Permit?

A Tree Removal & Replacement Permit is required for any tree:

- In an environmentally critical area or Tree Conservation Easement (TCE).
- Over six inches in diameter (outside of a critical area or TCE).
- That has been planted as a replacement tree for a previous permit.

To read the full regulations, please refer to Chapter 16.14 of our municipal code.

To learn more about the tree permit process, please visit our website at: https://www.cityoflfp.gov/163



Permits provide a city arborist with the opportunity to engage property owners:

- Encouraging retention of large trees
- Promoting tree health
- · Providing tree care information
- Monitoring canopy levels
- Ensuring trees are replaced



For more information, visit our website:

https://www.cityoflfp.com/107/Tree-Board

LAKE FOREST PARK Washington

his is a fantastic community; and we know you'll love it here. We care about our neighbors and respect the Urban Forest that surrounds us.

The Lake Forest Park Tree Board is here to help you get better acquainted with our **Urban Forest**



A city where people enjoy "the stately cedar, the majestic fir, and the homelike maple."



Get to Know Your Tree Board

THE TREE BOARD HAS THREE PRIMARY **RESPONSIBILITIES:**







For more information, visit our website: https://www.cityoflfp.com/107/Tree-Board




THE TREE BOARD HAS VACANCIES:



The city is seeking applications for vacant seats on the Lake Forest Park Tree Board. If you are interested in applying or have questions, please see our website and contact the Assistant Planner:

https://www.cityoflfp.gov/107/Tree-Board





Ole Hanson developed Lake Forest Park circa 1912 "with an eye on Nature." The same is true today.



Why are Tree Permits Important?


- Tree removal without subsequent tree replacement can cause potential water runoff problems for your property and neighboring properties.
- Permits can also help to avoid accidental cutting of trees on public property and in environmentally critical areas.



To learn more about the tree permit process, please visit our website at: https://www.cityoflfp.gov/163



Social Media: Instagram & Facebook











Get to Know Your Tree Board

THE TREE BOARD HAS THREE PRIMARY RESPONSIBILITIES:

- Educate and enlighten the community on tree-related issues
- Organize and facilitate the city's tree planting events and other public events involving trees
- Provide advice to City Council on policy and regulatory issues involving trees



For more information, visit our website: https://www.cityoflfp.com/107/Tree-Board



This is a fantastic community; and we know you'll love it here. We care about our neighbors and respect the Urban Forest that surrounds us.

The Lake Forest Park Tree Board is here to help you get better acquainted with our Urban Forest



A city where people enjoy "the stately cedar, the majestic fir, and the homelike maple."






For more information, visit our website: https://www.cityoflfp.com/107/Tree-Board



Why are Tree Permits Important?



 Permits can also help to avoid accidental cutting of trees on public property and in environmentally critical areas.



To learn more about the tree permit process, please visit our website at: https://www.cityoflfp.gov/163



The Tree Board has Vacancies

The city is seeking applications for vacant seats on the Lake Forest Park Tree Board. If you are interested in applying or have questions, please see our website and contact the Assistant Planner.

Board members should live within the Lake Forest Park city limits, but it is not required. Interest or background in urban forestry, horticulture, or habitat restoration are encouraged, but but required. Membership terms are for three years.

For more information, visit our website: https://www.cityoflfp.com/107/Tree-Board

Get to Know Your Tree Board

- Educate and enlighten the community on tree-related issues
- Organize and facilitate the city's tree planting events and other public events involving trees
- Provide advice to City Council on policy and regulatory issues involving trees

Getting to know your City's Arborist

- Helping the community to protect trees for future generations
- Managing ongoing challenges including: Civic construction projects
- Monitoring for drought, aging trees, disease
- Seasonal maintenance issues

Why are Tree Permits Important?

- Tree removal without subsequent tree replacement can cause potential water runoff problems for your property and neighboring properties.
- Permits can also help to avoid accidental cutting of trees on public property and in environmentally critical areas.
- Tree Permits let home owners easily remove problem trees with the provision that a new tree be replanted
- Permits help balance removals so that the neighborhood tree canopy remains balanced
- Permits provide a city arborist with the opportunity to engage property owners: Encouraging retention of large trees, Monitoring canopy levels, and Ensuring trees are replaced

To learn more about the tree permit process, please visit our website at:

https://www.cityoflfp.gov/163

KEY: Holiday eNews eNews DUE September, October, November 2022 Webinar **Blog Post** SlideShare Campaign Experiment Other **SUNDAY MONDAY TUESDAY WEDNESDAY THURSDAY FRIDAY** SATURDAY **SUNDAY SATURDAY MONDAY TUESDAY WEDNESDAY THURSDAY FRIDAY** 5 6 10 SUNDAY SATURDAY **MONDAY TUESDAY WEDNESDAY THURSDAY FRIDAY** 12 13 11 15 16 1) Impact of Heat <edit> 2) Leaves Rains Drains <edit > 3) Tree Concerns <edit > **SUNDAY MONDAY TUESDAY WEDNESDAY THURSDAY FRIDAY** SATURDAY 23 18 19 20 21 22 24

Section 8, ItemA.

			eNews DUE			
_						
SUNDAY	MONDAY	TUESDAY	WEDNESDAY	THURSDAY	FRIDAY	SATURDAY
25	26	27	28	29	30	
			eNews			

					CHARACTER
DATE	MESSAGE	LINK	CAMPAIGN	IMAGE	COUNT
	Recommendations for your neat-damaged trees and				
	shrubs				
	Go ahead and cut off dead flowers but try to resist the urge to			https://www.cityoflfp.com/A	
	remove partially dead leaves.	https://www.oregonlive.com/		reas/CivicSend/Assets/Uplo	
	Water deeply. Use drip or soaker hoses for more efficient water			ads/4953/s6b55e694adfb4	
	use; water is lost to evaporation when using an overhead	heat-damaged-plants-after-		d9e939dbd4df2a669ce637	
	sprinkler.	oregons-historic-heat-		684446943338567_optimiz	
September 2022	Apply mulch 2 to 3 inches deep.	wave.html	1) Impact of Heat <edit></edit>	<u>ed.jpg</u>	343
	It is time to check the storm drains and clear leaves from them				
	to keep the stormwater flowing.				
	Please do not create piles of leaves expecting the sweeper to				
	sweep them away. Use your yard waste tote so they can be				
	composed.				
	DO NOT rake or blow leaves and debris into the street,				
	ditchlines, or rights-of-way.				
	The sweeper cannot handle large piles of leaves. If the				
	sweeper encounters large piles of leaves, the driver must drive				
	around them, or the pile of leaves could damage the street				
	sweeper and the process may come to a screeching halt.				
	Municipal Code 16.25.025 makes it illegal to collect lawn				
	clippings, leaves or branches and discharge them into the path			https://www.cityoflfp.com/A	
	of surface water.			reas/CivicSend/Assets/Uplo	
	Never try to clear a storm drain or culvert if there is moving			ads/4953/240744b34a7446	
	water greater than knee deep, and always be wary of traffic			f4804c20f0a777a663_small	
October 2022	when working near a roadway.		2) Leaves Rains Drains <edit></edit>	_optimized.jpg	836
	As we enter the rainy/windy season, it can be natural to have				
	some concerns about trees on your property.				
	If a tree on your property is fully dead with no live foliage, you				
	can email a picture of the tree along with a description and your				
	address to the City Arborist for confirmation that the tree meets				
	the code definition of a dead tree.				
	If you'd like someone to come take a look at your trees and				
	provide an assessment of their health and/or structural				
	condition, the City recommends finding an ISA Certified			https://www.citvoflfp.com/A	
	Arborist.			reas/CivicSend/Assets/Uplo	
	If you'd like to remove any live trees on your property that have			ads/4953/0f149d9d245f40c	
	a DBH (diameter at 4.5 feet above ground) of 6 inches or more,			f8cb46ade2fd3024c_small_	
November 2022	then a tree removal permit application can be submitted		3) Tree Concerns <edit></edit>		709

KEY: Holiday eNews eNews DUE December 2022, January February 2023 Webinar **Blog Post** SlideShare Campaign Experiment Other **SUNDAY MONDAY TUESDAY WEDNESDAY THURSDAY FRIDAY SATURDAY SATURDAY SUNDAY MONDAY TUESDAY WEDNESDAY THURSDAY FRIDAY** 5 6 10 SATURDAY **SUNDAY MONDAY TUESDAY WEDNESDAY THURSDAY FRIDAY** 12 11 13 15 16 Dec: Christmas Tree Dilema Jan: Beaked Hazel Feb: Green Giants **SUNDAY MONDAY TUESDAY WEDNESDAY THURSDAY FRIDAY** SATURDAY 23 18 19 20 21 22

Section 8, ItemA.

			eNews DUE			
_						
SUNDAY	MONDAY	TUESDAY	WEDNESDAY	THURSDAY	FRIDAY	SATURDAY
25	26	27	28	29	30	
			eNews			

					CHARAC
DATE	MESSAGE	LINK	CAMPAIGN	IMAGE	COUNT
	Artificial?				
	Your volunteer Tree Board focuses on LFP's living forests. This time of year, though, we are often asked about Christmas trees, and which has a smaller environmental impact: buying a				
	real tree or an artificial one.				
	The bottom line: real! Real trees have a smaller carbon footprint and help fight climate change, and even though your				
	Christmas tree is cut down, you are actually supporting forests.				
	In the U.S., around 10 million artificial trees are purchased each season. Nearly 90 percent are shipped from China, resulting in				
	an increase in carbon emissions. Most are made of PVC, which pollutes across its entire lifespan, from production to end of life.				
	Most artificial trees are not recyclable and end up in landfills, where they release more greenhouse gases and leach				
	dangerous chemicals.				
	But shouldn't we avoid cutting down real trees? One of the best ways to protect forests is to use them-carefully. When our forests are sustainably managed, they can produce renewable				
	resources like Christmas trees and wood products. Christmas tree farms provide clean air and water, habitat for wildlife, and				
	erosion control. When these natural trees are harvested, there are more than 10 times as many left standing. For every tree				
	purchased, farmers plant 1-3 seedlings in its place. Plus, most of the 15,000 Christmas tree farms across the U.S. are family-				
	owned, so when you buy a real Christmas tree, you support local economies and contribute to an industry that provides				
	100,000+ jobs.				
	The caveat: try to ensure your real tree comes from a local, sustainable farm that doesn't use pesticides, and make sure it				
	is responsibly composted after the holidays.				
	Maybe you are considering a living tree to plant in your yard after the holidays. If you keep it inside for less than a week, then plant it promptly, it should survive and become a valuable				
	addition to our city's tree canopy. Any longer inside than that, though, and they rarely survive the transition. A small-scale			Treelotstockimage.jpg	
	alternative is a potted Norfolk Island Pine, which is happy as a year-round houseplant.			Or	
	If you still prefer the convenience of an artificial tree, the key is			https://www.cityoflfp.com/A reas/CivicSend/Assets/Uplo	
	to buy one that's high quality, so it will last. An artificial tree must be in circulation for at least 8 yearsbut ideally 20to		1) Look for an Evergreen is Dec	ads/4953/s15bf426d7e934 dd7a9b5c37797edee06_sm	
December 2022	negate its carbon footprint. Or go vintage! Scout the many used Meet the Beaked Hazel – Corylus cornuta		19,2022 <edit></edit>	all_optimized.jpg	2516
	Lake Forest Park is proud of its extensive trees, but a forest				
	includes much more than its canopy trees. Our attention is sometimes drawn lower in the forest to subcanopy trees that				
	are also a critical part of our urban forest community.				
	The beaked hazel (<i>Corylus cornuta</i>) is the earliest native flowering plant in our forests, appearing as a tawny yellow haze, and often seen flowering best where it grows, along the				
	edge of a forest. Flowering in January and February at low elevations and later in the spring as you go up in elevation.				
	Like other members of the birch family, beaked hazel flowers				
	before the leaves emerge in the spring. This is not by accident; they are all wind pollinated and dense foliage inhibits both free				
	dispersal and capture of pollen. Wind-pollinated trees typically separate the male stamens and female pistils into separate				
	flowers. In fact, wind pollination is so inefficient that plants with separate male and female flowers allocate more resources to				
	pollen production by having more male flowers to make up for all of the wasted pollen.				
	From a distance, male flowers, which are arranged in pendant "catkins" (a spike of unisexual, apetalous flowers having scaly,				
	usually deciduous bracts, as of a willow or birch) create the yellow aura that signals the presence of a hazel in our early				
	spring woods. While not visible from a distance, it is worth taking a closer look to see the tiny, brilliant, red female flowers				
	arranged individually near the tips of stems. The bright red structures are not the petals we normally think of as the colorful				
	part of a flower, but are the stigma, where pollen is deposited. On plants pollinated by animals, the stigma is small and				
	unobtrusive; but, in wind-pollinated plants, it is large and branched to capture the pollen as it blows by on the wind,				
	instead of having the pollen delivered to it. Since it does not need to attract animal pollinators, it is not known why they are such a brilliant red; maybe it's to attract our attention!				
	In birches and alders, the female flowers are also organized in				
	catkins. Why not in hazel? Birches and alders produce lots of small, winged seeds that are dispersed by the wind the				
	following fall, but hazelnuts are heavy and a small branch tip would not be able to support a bunch of them. Like the				
	commercial European hazelnut, ours is edible, but they never seem to produce in large quantities and we are unlikely to beat				
	the squirrels to the ones they do produce. In late summer, we can see big gray squirrels (introduced from eastern North			https://www.cityoflfp.com/A reas/CivicSend/Assets/Uplo	
2022	America) foraging in hazels for the still-green nutssmall branches waving to and fro as the squirrels move from nut to nut.		200-1-111-1-1700	ads/4953/sbad48ff64d0741 3ea9d8e8851ad55ef2_smal	2915
January 2023	Groom Granto		2) Beaked Hazel (TBD)	L_optimized.jpg	2913
	Here in the Pacific Northwest, we share our home with very diverse botanical giants. Most conifers are trees, although a				
	handful are shrubs. An iconic trait of conifers is their reproductive cone, often composed of overlapping scales: large				
	and woody in western white pine, medium size with protruding "mouse tail" bracts in Douglas fir, and very small, upright ovoid				
	cones with few scales on western red cedar. With many shaped like the typical Christmas tree, conifers have produced the world's tallest, thickest, and oldest specimens. Every type of				
	world's tallest, infecest, and oldest specimens. Every type of coniferous tree represented in the Northwest finds its largest and often longest-lived individuals here.				
	We are lucky in Lake Forest Park to have such a botanical				
	backdrop for our daily lives. How often do we pause to take stock of such a glorious natural legacy? Traveling to my family				
	home in the Midwest, I feel unnaturally exposed within what seems the stunted stands of native deciduous trees. When I				
	return to Lake Forest Park, it is like emerging from a tunnel into a dazzling dreamworld of green and verdant growth reaching toward the sky.				
	toward the sky. Visitors often ask if our trees are remnant old growth and are				
	stunned when I tell them our trees are youngsters, mostly 50 – 100 years in age. With care and preservation, they may live				
	several hundred more years. More surprisingly still, big Douglas firs are outlived by Western hemlock and Western red			https://www.cityoflfp.com/A reas/CivicSend/Assets/Uplo	
	cedar—some living perhaps 1,000 years or more. The grand and beautiful coniferous forests of the Northwest are a			ads/4953/s33bdb8d2dc184 6abb5e001f311413eac_sm	
February 2023	botanical treasure of immeasurable importance to the health	I	3) Green Giants (TBD)	all_optimized.jpg	1658

Section 8, ItemA.

25

DATE	CAMPAIGN	MESSAGE	LINK
Month of March	Gardening, Nature, and Ecology Books Month		
March 7th	Plant Power Day		
March 11th & Sept 26th	Johnny Appleseed Day		
March 20th	International Earth Day & Spring Equinox		
March 22nd	National Agriculture Day		
Month of April	National Garden Month		
April 13th	International Plant Appreciation Day		
April 14th	National Gardening Day		
April 22nd	Earth Day		
Last Friday in April	Arbor Day		
Month of May	Gardening for Wildlife Month		
2nd week in May	National Public Gardens Week		
May 3rd	Garden Mediation Day		
May 6th	National Public Gardens Day		
May 16th	Love a Tree Day		
May 19th	Plant Something Day		
3rd Saturday in May	Plant a Lemon Tree Day		
1st week of June	National Gardening Week		
June 5th	World Environment Day		
Month of July	National Outdoor Month		
December 19th	Look for an Evergreen Day		
Summer	IVY OUT Events		

CONTENT			
TYPE	TITLE	INTERESTING SNIPPET	IMAGE
		No and a triangle desired time of the control of th	
		We are entering the driest time of the year - so unless your garden only has established drought-tolerant plants, summer watering is a necessity. It is generally better to water deeply and infrequently. Moisten the whole root zone	
		and let the soil dry before watering again. In hot weather, garden beds and lawns (unless you let your grass go dormant) need about an inch of water per week.	https://www.cityo flfp.com/Areas/Ci
			vicSend/Assets/U
	Summer Garden	Make every drop count by mulching, selecting drought-tolerant plants, using soaker hoses and water timers, and watering only in the early morning or evening to reduce evaporation.	ploads/4953/220 a7d3fb0aa4b4c9
	Watering		b42a372b48be6f
August	Advice	More information is available <u>here</u> .	9_optimized.jpg
		From the Tree Board: Illustrated Tree Walks in Lake Forest Park	
		This is part of a series of occasional columns on the trees of Lake Forest Park. The LFP Tree Board exists to advise the City on policies pertaining to trees, to facilitate tree planting events, and to provide outreach and education to the community.	
		Citizens of Lake Forest Park are rightfully proud of its extensive tree canopy and all the benefits that provides. Sometimes it is hard to see the trees, for the forest! Our urban forest includes an incredible diversity of trees in City parks, street rights-of-way, and especially in residents' gardens throughout the City.	
		It has been nearly 15 years since LFP resident, David Hepp, and the Urban Forest Task Force created the wonderful booklet <i>Tree Walks in Lake Forest Park</i> . The booklet describes four walks through different neighborhoods in Lake	https://www.cityo flfp.com/Areas/Ci
		Forest Park, pointing out and identifying noteworthy trees, both native and ornamental, encountered along each route.	vicSend/Assets/U
			d9fbfeb1d48869
		Because neighborhoods can change—trees grow or die, and new ones are planted—the LFP Tree Board, with	91fea623d7b6e0
		assistance from David Hepp, is revising the Tree Walks. In addition to updating the original descriptions by removing	8637920288784
	Illustrated	trees that have died and including new ones that caught our attention, photos of many of the trees are included to	646699_small_op
July	Tree Walks	help users pick out the trees noted in the text. The new Illustrated Tree Walks booklet is available online and can be	timized.jpg

		Spring Gardening Spring gardening is right around the corner! This year, try gardening without all the chemicals. Get free one-on-one garden advice from www.gardenhotline.org . Learn more about how to use Natural Yard Care methods at www.naturalyardcare.org .	https://www.cityo flfp.com/Areas/Ci vicSend/Assets/U ploads/4953/a4b 5ecef605d4e2eb 14b20f136281eb
	Spring		5_small_optimize
February	Gardening		<u>d.jpg</u>

The omisimas free bhemma. Near of Artificial:

Submitted by the LFP Tree Board

Your volunteer Tree Board focuses on LFP's living forests. This time of year, though, we are often asked about Christmas trees, and which has a smaller environmental impact: buying a real tree or an artificial one.

The bottom line: real! Real trees have a smaller carbon footprint and help fight climate change, and even though your Christmas tree is cut down, you are actually supporting forests.

In the U.S., around 10 million artificial trees are purchased each season. Nearly 90 percent are shipped from China, resulting in an increase in carbon emissions. Most are made of PVC, which pollutes across its entire lifespan, from production to end of life. Most artificial trees are not recyclable and end up in landfills, where they release more greenhouse gases and leach dangerous chemicals.

But shouldn't we avoid cutting down real trees? One of the best ways to protect forests is to use them—carefully. When our forests are sustainably managed, they can produce renewable resources like Christmas trees and wood products. Christmas tree farms provide clean air and water, habitat for wildlife, and erosion control. When these natural trees are harvested, there are more than 10 times as many left standing. For every tree purchased, farmers plant 1-3 seedlings in its place. Plus, most of the 15,000 Christmas tree farms across the U.S. are family-owned, so when you buy a real Christmas tree, you support local economies and contribute to an industry that provides 100,000+ jobs.

The caveat: try to ensure your real tree comes from a local, sustainable farm that doesn't use pesticides, and make sure it is responsibly composted after the holidays.

Maybe you are considering a living tree to plant in your yard after the holidays. If you keep it inside for less than a week, then plant it promptly, it should survive and become a valuable addition to our city's tree canopy. Any longer inside than that, though, and they rarely survive the transition. A small-scale alternative is a potted Norfolk Island Pine, which is happy as a year-round houseplant.

The Christmas

December Tree Dilema

If you still prefer the convenience of an artificial tree, the key is to buy one that's high quality, so it will last. An artificial tree must be in circulation for at least 8 years--but ideally 20--to negate its carbon footprint. Or go vintage! Scout the

https://www.cityo flfp.com/Areas/Ci vicSend/Assets/U ploads/4953/s15b f426d7e934dd7a 9b5c37797edee0 6_small_optimize d.ipg

Impact of June "Heat Dome" on LFP Trees

Submitted by Richard Olmstead, LFP Tree Board

We will all have stories for years to come of the infamous Pacific Northwest "heat dome" that set and then broke temperature records throughout the region for three days in a row, culminating in 108°F in Seattle on June 28. It was too much figuratively, and, sadly, literally, for many people in our region. But what about the trees that shape the identity of Lake FOREST Park?

First, let's consider the good news. The temperature in my woodland garden rose to 100°F. That's hot, but the extensive forest canopy that our city is blessed with has a buffering effect on temperature, so no matter how hot it seemed, we had it better than many communities in our area.

Over the next few days, however, the effect of that bright sun and heat on our trees and vegetation became visible, with browning leaves evident on a broad range of trees and shrubs, especially where exposed to the sun during the hottest parts of the day. The visible damage was frightening to anyone who loves trees, but how bad really was it?

Long-lived trees and shrubs have evolved to survive a range of conditions in their native environments, but the recordbreaking heat they experienced this year created stresses beyond the normal range of variation. It is informative to see how they responded.

The combination of bright sun and hot temperatures creates a series of stresses for plants, which of course are fixed in place and can't retreat into air-conditioned homes. While sun and warm temperatures are necessary for their photosynthesis and growth, excess of both can create a demand for water that a plant cannot keep pace with. This causes the physiology within leaves to malfunction and their cells to die. The immediate result can be patches of dead leaves on many trees, but it can also have a more measured effect in prioritizing resources—water--within the tree to sacrifice some leaves, while allocating available resources to the leaves most needed for survival.

In some cases, the damage was restricted to the leaves immediately affected by the heat and sun, leaving patches of dead leaves in the most exposed positions at the tips of branches. In other cases, trees were able to cut off water to older, less efficient leaves while maintaining younger leaves on branch tips. In my garden, I've seen examples of both patterns of damage. Incense cedar and salal suffered loss of leaves on branch tips that were most exposed, while noble fir and Wilson's magnolia lost the older leaves, keeping only the youngest leaves on branch tips. Though unsightly this summer, they will survive and look fine next year.

I'm also confident that trees in my garden will survive because I was able to provide supplemental water this summer. Of greater concern than the heat wave in June is the fact that Seattle received the least spring and summer rainfall in 2021 since records have been kept. This prolonged drought is much more likely to result in tree mortality than the record-breaking but short-term "heat dome" of June.

Recommendations for your heat-damaged trees and shrubs (with info from *The Oregonian*)

Go ahead and cut off dead flowers but try to resist the urge to remove partially dead leaves. Leaves, dead or alive,

August	Beat the	Beat the Heat with Natural Yard Care Sick of the heat yet? Well, we'll need to get used to it, because hot, dry summers are going to continue to be the norm here in the Puget Sound region. For gardeners, this may mean having to re-imagine what your garden looks like and how you tend to it, in order to adapt to a changing climate. Luckily, following these five simple steps to natural yard care can help by following this link: https://pugetsoundstartshere.today/2021/07/26/beat-the-heat-with-natural-yard-care/	https://www.cityo flfp.com/Areas/Ci vicSend/Assets/U ploads/4953/70b a6788b4934a198 24c8d3b994b104 5_small_optimize d.jpg
		Green Giants Timothy Hohn, LFP Tree Board	
		Here in the Pacific Northwest, we share our home with very diverse botanical giants. Most conifers are trees, although a handful are shrubs. An iconic trait of conifers is their reproductive cone, often composed of overlapping scales: large and woody in western white pine, medium size with protruding "mouse tail" bracts in Douglas fir, and very small, upright ovoid cones with few scales on western red cedar. With many shaped like the typical Christmas tree, conifers have produced the world's tallest, thickest, and oldest specimens. Every type of coniferous tree represented in the Northwest finds its largest and often longest-lived individuals here.	
February		We are lucky in Lake Forest Park to have such a botanical backdrop for our daily lives. How often do we pause to take stock of such a glorious natural legacy? Traveling to my family home in the Midwest, I feel unnaturally exposed within what seems the stunted stands of native deciduous trees. When I return to Lake Forest Park, it is like emerging from a tunnel into a dazzling dreamworld of green and verdant growth reaching toward the sky. Visitors often ask if our trees are remnant old growth and are stunned when I tell them our trees are youngsters, mostly 50 – 100 years in age. With care and preservation, they may live several hundred more years. More surprisingly still, big Douglas firs are outlived by Western hemlock and Western red cedar—some living perhaps 1,000 years or more. The grand and beautiful coniferous forests of the Northwest are a botanical treasure of immeasurable importance to the health and	ploads/4953/s33b

This is the first of occasional columns on the trees of Lake Forest Park. The LFP Tree Board exists to advise the City on policies pertaining to trees, to facilitate tree planting events, and to provide outreach and education to the community.

Lake Forest Park is proud of its extensive trees, but a forest includes much more than its canopy trees. Our attention is sometimes drawn lower in the forest to subcanopy trees that are also a critical part of our urban forest community.

The **beaked hazel** (*Corylus cornuta*) is the earliest native flowering plant in our forests, appearing as a tawny yellow haze, and often seen flowering best where it grows, along the edge of a forest. Flowering in January and February at low elevations and later in the spring as you go up in elevation.

Like other members of the birch family, beaked hazel flowers before the leaves emerge in the spring. This is not by accident; they are all wind pollinated and dense foliage inhibits both free dispersal and capture of pollen. Windpollinated trees typically separate the male stamens and female pistils into separate flowers. In fact, wind pollination is so inefficient that plants with separate male and female flowers allocate more resources to pollen production by having more male flowers to make up for all of the wasted pollen.

From a distance, male flowers, which are arranged in pendant "catkins" (a spike of unisexual, apetalous flowers having scaly, usually deciduous bracts, as of a willow or birch) create the yellow aura that signals the presence of a hazel in our early spring woods. While not visible from a distance, it is worth taking a closer look to see the tiny, brilliant, red female flowers arranged individually near the tips of stems. The bright red structures are not the petals we normally think of as the colorful part of a flower, but are the stigma, where pollen is deposited. On plants pollinated by animals, the stigma is small and unobtrusive; but, in wind-pollinated plants, it is large and branched to capture the pollen as it blows by on the wind, instead of having the pollen delivered to it. Since it does not need to attract animal pollinators, it is not known why they are such a brilliant red; maybe it's to attract our attention!

In birches and alders, the female flowers are also organized in catkins. Why not in hazel? Birches and alders produce lots of small, winged seeds that are dispersed by the wind the following fall, but hazelnuts are heavy and a small branch tip would not be able to support a bunch of them. Like the commercial European hazelnut, ours is edible, but they never seem to produce in large quantities and we are unlikely to beat the squirrels to the ones they do produce. In late summer, we can see big gray squirrels (introduced from eastern North America) foraging in hazels for the still-green nuts--small branches waving to and fro as the squirrels move from nut to nut.

https://www.cityo flfp.com/Areas/Ci vicSend/Assets/U ploads/4953/sbad 48ff64d07413ea 9d8e8851ad55ef 2_small_optimize d.jpg

January Hazel

Timothy Hohn, Chair, LFP Tree Board

Something incredible is happening underground in our community forest. Scientists have discovered that forest trees and understory plants are all linked together in a "wood-wide web," a mutually supportive network of fungal threads and plant roots. The network is anchored by large "mother" trees that act as network hubs, doling out surplus resources to smaller trees and plants. In effect, these networked groves of monarch trees and their lesser green mortals are one symbiotic association.

With this in mind, I met with the Lake Forest Park City Council on behalf of the Tree Board to pass along Board recommendations on how the city code might better support our community forest wood-wide web. These recommendations included code references to tree "groves" and how they are defined, as well as new dimensions for important "exceptional" trees—what we now know are grove "mother" trees—that form the foundation of our community forest and its wood-wide web. We also discussed prioritizing native trees as replacements for those that are removed, in order to retain the ecological integrity of our community forest. The City Council is likely to ask the Planning Commission to review these suggestions early next year. To view the recording of my presentation and the Council's discussion, click here.

https://www.cityo flfp.com/Areas/Ci vicSend/Assets/U ploads/4953/s8ce 4e0d091034f23a 74ad278fc5eabe 8_small_optimize d.jpg

December Our Forest

As we move into the rainy season and leaves start to fall, the Public Works Department starts turning its attention to the City's drainage system. It is time to check the storm drains and clear leaves from them to keep the stormwater flowing. Street sweeping is a component of the drainage system maintenance and is perhaps the most visible to the citizens. The City is encouraging residents to clear any blocked storm drains that are adjacent to or in front of their property keeping in mind:

Please do not create piles of leaves expecting the sweeper to sweep them away. Use your yard waste tote so they can be composed.

DO NOT rake or blow leaves and debris into the street, ditchlines, or rights-of-way.

The sweeper cannot handle large piles of leaves. If the sweeper encounters large piles of leaves, the driver must drive around them, or the pile of leaves could damage the street sweeper and the process may come to a screeching halt.

Municipal Code 16.25.025 makes it illegal to collect lawn clippings, leaves or branches and discharge them into the path of surface water.

Never try to clear a storm drain or culvert if there is moving water greater than knee deep, and always be wary of traffic when working near a roadway.

Wet leaves are surprisingly heavy, so be careful not to overexert yourself. Debris from storm drains should be placed in yard waste containers. If flooding is severe, or you find evidence of dumping, please call 206-368-5440.

What can you do with leaves?

Turn your leaves into beneficial compost. Autumn leaves are a great source of high-carbon material for your compost pile. Alternate layers of shredded leaves with the other materials you normally add to your compost pile (scraps from fruits and veggies, grass clippings, weeds, etc.) and let it sit over the winter. Whenever you think about it, aerate or turn the pile. Your compost will be ready to use by spring.

Shred leaves and **use them as mulch** on vegetable gardens and flower beds, around trees and shrubs, and in containers. Just add a 2- to 3-inch layer of shredded leaves to the beds, keeping the mulch from directly touching the stems and trunks of the plants. The mulch retains moisture in the soil, stays cool, and limits weed seed germination. Use a mulching mower to shred your leaves once a week until they have all fallen, then **"leave" them on your yard.** While the leaves break down during winter, they'll shade your soil and provide it with nutrients, which means fewer weeds to deal with in spring. **Hoard them!** Save a bag or two of leaves in your garage over the winter. In spring, adding that brown material to

https://www.cityo flfp.com/Areas/Ci vicSend/Assets/U ploads/4953/240 744b34a7446f48 04c20f0a777a66 3_small_optimize d.ipg

Leaves,

October Rains, Drains your compost pile makes your compost just right for the season.

15

	As we enter the rainy/windy season, it can be natural to have some concerns about trees on your property. The Planning Department staff is here to help offer some refreshers on best practices and permit requirements. If a tree on your property is fully dead with no live foliage, you can email a picture of the tree along with a description and your address to the City Arborist for confirmation that the tree meets the code definition of a dead tree. Once you receive confirmation from the City Arborist, then you will be allowed to remove the dead tree without a permit. If you'd like someone to come take a look at your trees and provide an assessment of their health and/or structural condition, the City recommends finding an ISA Certified Arborist. A Certified Arborist can evaluate your trees of concern and provide helpful information about the health of your trees and if any problems can be mitigated through	
	pruning or other measures. If you'd like to remove any live trees on your property that have a DBH (diameter at 4.5 feet above ground) of 6 inches or more, then a tree removal permit application can be sent via email to either the City Arborist or the Assistant Planner and payments can be made by check. Checks should be made out to "City of Lake Forest Park" in the amount of \$76.13 (application fee) and can be submitted: By mail to City Hall, 17425 Ballinger Way NE, Lake Forest Park, 98155; or Dropped off at City Hall in a letter-sized envelope in the silver payment drop box (located on the west side of the building next to the King County elections box). Once we receive the application and payment, we will contact you to discuss next steps. If it becomes apparent that a live tree needs to be removed immediately and is an emergency, the City code does have provisions for emergency actions. We recommend contacting the City Arborist and the Assistant Planner for	https://www.cityo flfp.com/Areas/Ci vicSend/Assets/U ploads/4953/0f14 9d9d245f40cf8cb 46ade2fd3024c_s
October	guidance as soon as you become aware that a tree may need to be removed as an emergency.	mall_optimized.jp
	Wondering how to remove ivy safely? The King County Noxious Weed Board has tips on how to identify ivy (page 2) and recommend manual removal practices (page 3). You can also click here to learn about long-term maintenance ideas and why ivy removal is important for our urban ecosystems.	https://www.cityo flfp.com/Areas/Ci vicSend/Assets/U
July lvy C	Please note that removal of invasive plants is considered to be exempt from permitting requirements, unless the invasive plants are located within a steep slope hazard area or its buffer. If you are unsure whether the proposed removal area is located within a steep slope hazard area or have other questions, please contact the Assistant Planner for more information.	ploads/4953/ca18 1fbac4964720a8 30850259f1c467 _small_optimized. jpg

		martir bay to / a variar are variar.	
		Did you know there is no filtration system between your outdoor drains and our streams, Lake Washington, or Puget Sound? Where you wash your car, not picking up dog poop, leaky vehicles, improper application of yard care products, and more all contribute to the water quality in Lake Forest Park and around the Sound.	
		Now is a good time and reminder to do a little something extra to help the earth and our LFP environment. A few examples of things you could do at home are:	
		Wash your vehicles at a commercial car wash where they treat the dirty water properly.	
		Pick up your dog's poop—remember to always have a bag with you and put it in the trash. Going on a walk? Pick up	
		the extra pile you pass by.	https://www.city
		Fix that leaking car to improve its longevity and our environment.	flfp.com/Areas/C
		Properly apply yard care products and learn some natural yard care techniques. For advice check out the Garden	vicSend/Assets/l
		Hotline or call them at 206-633-0224.	ploads/4953/scc
		Use and dispose of chemicals properly. King County provides some great options and resources.	6a232dc3248df
		Maintain your septic system, don't let that waste water into our groundwater!	70fee1ecd6878
		Click on the link for more information on these topics or other options of what you can do to help the environment, We	4 small optimize
March	Earth Day	know it's not easy, but do what you can to help water quality and bring salmon back to our streams, lakes, and	d.jpg
	_	Earth Smart Green Fair	
		Learn to save the planet at Lake Forest Park's 17th Annual <i>Earth Smart Green Fair</i> ! All are welcome and invited to	
		attend – and it's free! The fair is hosted at Third Place Commons (top floor) from 10:00 a.m. to 2:00 p.m., March 21,	
		2020. Bring your friends, neighbors, and family and come on down for some fun, learn something new, and take	https://www.city
		home some giveaways!	flfp.com/Areas/C
			vicSend/Assets/l
		Local organizations and businesses will be there to answer your questions regarding composting, recycling, reducing	ploads/4953/sb6
		waste, natural yard care, water conservation, environmentally-safe products, and more—all for free!	<u>b69c0a3a946ef8</u>
			4a4222e299122
	Earth Smart	View our <u>Facebook page</u> for more information!	5_small_optimize
February	Green Fair		<u>d.jpg</u>





GETTING TO KNOW YOUR CITY'S ARBORIST



Helping the community to protect trees for future generations.

Managing ongoing challenges including:

- Civic construction projects
- Drought, aging trees, disease
- Seasonal maintanance issues


"Ensure tree safety, value, and beauty".



"Maintain, expand, enhance our urban forest"







https://www.cityoflfp.com/107/Tree-Board





Community Engagement

Permits provide a city arborist with the opportunity to engage property owners:

- Encouraging retention of large trees
- Promoting tree health
- Providing tree care information
- Monitoring canopy levels
- Ensuring trees are replaced



For more information, visit our website:

https://www.cityoflfp.com/107/Tree-Board

LAKE FOREST DAD Washington

Section 8. ItemB.

This is a fantastic community; and \divideontimes we know you'll love it here. We care about our neighbors and respect the Urban Forest that surrounds us.



Tree Permits let home owners easily remove problem trees with the provision that a new tree be replanted



Permits help balance removals so that the neighborhood tree canopy remains balanced

Do I need to submit a Tree Permit?

A Tree Removal & Replacement Permit is required for any tree:

- In an environmentally critical area or Tree Conservation Easement (TCE).
- Over six inches in diameter (outside of a critical area or TCE).
- That has been planted as a replacement tree for a previous permit.



To read the full regulations, * please refer to Chapter 16.14 of our municipal code.

Section 8. ItemB.

LAKE FOREST PARK Washington

This is a fantastic community; and **
we know you'll love it here. We
care about our neighbors and
respect the Urban Forest that
surrounds us.



Tree Permits let home owners easily remove problem trees with the provision that a new tree be replanted



Permits help balance removals so that the neighborhood tree canopy remains balanced

To learn more about the tree permit process, please visit our website at: https://www.cityoflfp.gov/163

Section 8. ItemB.





A Tree Removal & Replacement Permit is required for any tree:

- In an environmentally critical area or Tree Conservation Easement (TCE).
- Over six inches in diameter (outside of a critical area or TCE).
- That has been planted as a replacement tree for a previous permit.



To learn more about the tree permit process, please visit our website at: https://www.cityoflfp.gov/163









GETTING TO KNOW YOUR CITY'S ARBORIST




Helping the community to protect trees for future generations.

Managing ongoing challenges including:

- Civic construction projects
- Drought, aging trees, disease
- Seasonal maintanance issues

"Ensure tree safety, value, and beauty".







Community Engagement

Permits provide a city arborist with the opportunity to engage property owners:

- Encouraging retention of large trees
- Promoting tree health
- Providing tree care information
- Monitoring canopy levels
- Ensuring trees are replaced




For more information, visit our websith https://www.cityoflfp.com/107/Tree-Boa



CITY OF LAKE FOREST PARK

URBAN FOREST ECOSYSTEM SERVICES AND VALUES REPORT

DECEMBER 2023



Acknowledgements

DCG/Watershed Contributors

Kim Frappier, Project Manager & Author Sam Payne, Data Analyst & Author Nathan Burroughs, GIS Analyst Lars Freeman-Wood, Lead Field Researcher Sage Yuasa, Field Researcher Evan Earhart, Field Researcher Drew Foster, Field Researcher & Analyst Kenny Booth, Principal

City of Lake Forest Park

Phillip Hill, ICMA-CM, City Administrator
Councilmember Larry Goldman, City Council
Steve Bennett, Planning Director
Hannah Swanson, City Arborist
Richard Olmstead, Tree Board Chair
Doug Sprugel, Tree Board
Marty Byrne, Tree Board
Sandra LeVar, Tree Board
Mark Phillips, Tree Board

DICIG WATERSHED





We wish to express our gratitude to the i-Tree project team for their invaluable contribution in developing methodologies and software which facilitated the analysis in this study. Contributors to the i-Tree project are listed as the U.S. Forest Service, Davey Tree Expert Company, The Arbor Day Foundation, Society of Municipal Arborists, International Society of Arboriculture, and Casey Trees.

Project funding is from the City of Lake Forest Park.

Uncredited photos by Sam Payne, or other DCG/Watershed project staff.





Summary

This report presents a comprehensive evaluation of Lake Forest Park's urban and community forest through an i-Tree Eco plot sample inventory. Utilizing plot data obtained in 2022 and 2023, the i-Tree Eco model provides an assessment of urban forest health, structure, and threats as well as the ecosystem services and values trees provide the community. In addition to the i-Tree analysis, this study compared tree canopy height using LiDAR to better understand the distribution of various canopy heights in the City's tree population. The following summarizes key findings from this research effort:

- There are a total of 297,100 trees estimated to be in Lake Forest Park with a mean density of 129 trees per acre (TPA).
- Canopy cover is estimated at 50.6%, a level similar to prior studies.
- The most common tree species are Douglas-fir (16%), bigleaf maple (11%), western red cedar (9%), cherry laurel (8%), bitter cherry (6%), and English holly (6%). Of all trees, 63% are native to Washington.
- Less than 1% of trees are designated as noxious weeds in King County, however, 19% are listed as weeds of concern. The most abundant weeds of concern are cherry laurel, English holly, and bird cherry.
- The age classification of trees trends youthful, with an abundance of smaller trees that will eventually replace the aging canopy.
- Leaf area density in the Large Residential stratum (parcels >¹/₄ acre) is 3 times greater than the Small Residential stratum (parcels ≤¹/₄ acre), and 7 times greater than the Town Center stratum.
- The Lake Forest Park urban forest provides benefits valued at \$4.1 million annually for removing pollution, reducing runoff, sequestering carbon, and lowering energy usage.
- Carbon storage of the total urban forest is valued at \$16 million, and the replacement value is estimated at \$531 million.
- Of the 53 pests and pathogens that i-Tree assessed, 15 are present in King County. The economic impacts of these species are evaluated for each tree species and pest species.
- The canopy height model indicates that the proportion of tall trees, those greater than 135 feet in height, have increased by 21% from 2016 to 2021. The proportion of the tallest trees, those greater than 165 feet increased by 86% during this period, albeit accounting for less than 1% of the total tree population.



Table of Contents

INTRODUCTION 2				
	PROJECT BACKGROUND AND OBJECTIVES	1		
	SUMMARY OF URBAN FOREST BENEFITS	3		
M	METHODS			
	I-TREE STUDY DESIGN	4		
	LIMITATIONS AND ASSUMPTIONS	5		
	CANOPY HEIGHT MODEL	6		
RE	SULTS	7		
	TREE CHARACTERISTICS OF THE URBAN FOREST	7		
	URBAN FOREST COVER AND LEAF AREA	9		
	AIR POLLUTION	11		
	CARBON SEQUESTRATION AND STORAGE	12		
	SURFACE WATER RUNOFF	13		
	TREE BENEFITS SUMMARY	14		
	PESTS AND PATHOGENS	15		
	CANOPY HEIGHT MODEL RESULTS	18		
CANOPY HEIGHT MODEL RESULTS DISCUSSION		19		
	CLIMATE ADAPTATION AND RESILIENCE	19		
	PROTECTION OF SIGNIFICANT & LARGE DIAMETER TREES	20		
	INVASIVE SPECIES MANAGEMENT	21		
	ADDITIONAL CONSIDERATIONS	21		
RE	FERENCES	23		

APPENDICES

APPENDIX I. I-TREE ECO MODEL AND FIELD MEASUREMENTS

APPENDIX II. RELATIVE TREE EFFECTS

APPENDIX III. COMPARISON OF URBAN FORESTS

APPENDIX IV. GENERAL RECOMMENDATIONS FOR AIR QUALITY IMPROVEMENT

 $\label{eq:continuous} \textbf{APPENDIX V. INVASIVE SPECIES OF THE URBAN FOREST}$

APPENDIX VI. POTENTIAL RISK OF PESTS



Introduction

Lake Forest Park's urban and community forest consists of street trees, forested parks and open spaces, as well as trees on private residential, commercial, and industrial properties. These urban forest resources provide numerous ecosystem services, public health, and economic benefits to the people who live, work and recreate here. Jurisdictions across King County and the State of Washington are faced with the need to support smart growth and development, environmental and climate sustainability, change resilience. Protecting green infrastructure such as tree canopies critical to addressing these public and environmental health issues while ensuring the livability of Lake Forest Park. The first critical step to stewarding and managing this natural resource is understanding what we have.

The City of Lake Forest Park program has invested in tree inventories, canopy cover modeling, and studies investigating urban forest structure and values to guide the urban forestry program. This data has been used to inform and guide management actions, policies, municipal code

updates, budget development, and identify additional analysis needs. To date, the City has developed the following urban forest analysis and management plans:

- 2005 and 2016 Canopy Analyses (LiDAR based studies)
- 2011 Urban Forest Effects and Values (i-Tree Eco Analysis)
- 2010 Community Forest Management Plan

Project Background and Objectives

To build from the previous i-Tree Eco study published in 2011, the City contracted with DCG/Watershed in 2022 to conduct a follow-up survey to assess Lake Forest Park's community forest 10 years later. This analysis was first conducted in 2011 by the City arborist and Lake Forest Park Tree Board, with community volunteers participating in plot data collection.

The primary objectives of this 2022-2023 i-Tree study are to characterize urban forest structure and composition by collecting data on tree size, species, and health conditions. This data, along with other site level information within the specific study areas is then used to calculate the environmental and economic benefits at a city-scale. For a summary of environmental and economic benefits provided by urban forests, see page 3.

Studying the structure and composition of the urban forest through the i-Tree analysis, provides us with a more detailed understanding of Lake Forest Park's city-wide tree canopy, which were conducted in 2005 and again in 2016 using LiDAR analysis.

Lake Forest Park's urban tree canopy covered 43% of the total City area in 2004, which became a baseline for forest management goals established in the City's 2010 Community Forest Management Plan. To reflect the diverse landscapes and development regulations within Lake Forest Park, canopy cover goals were established by land use types to be 50% in suburban residential areas (lots >1/4 acres), 35% in suburban residential areas (lots <1/4 acres), and 15% in business districts. These were informed in-part by benchmarks recommended by American Forests. By 2016, total urban forest canopy increased to 50% according to a study (Elm




Figure 1. Lake Forest Park Tree Canopy Cover in 2016, reproduced from Elm (2016).

2016). This is consistent with recent analysis from i-Tree Landscape using high resolution data from 2017 which resulted in a canopy cover of 48%.

Urban forest structure is defined as the horizontal and vertical arrangement of trees, shrubs, and other plants, and their underlaying abiotic environments, and is relevant to management because the physical arrangement in three-dimensional space influences the functions and ecosystem services provided by a forest. Composition refers to tree or other plant species that make up a forest.

Summary of Urban Forest Benefits



Pollution Abatement: Urban forests serve as natural filters which improve water quality and air quality by trapping, absorbing, and transforming pollutants and excess nutrients, resulting in public health benefits, lower illness rates, and safeguarding ecosystems.



Shade and Cooling: Cities and metropolitan areas experience greater temperatures due to land use changes which alter the energy budget in an urban setting, known as the urban heat island effect. Through shading and evapotranspiration, urban forests mitigate the heat island effect through shading and cooling which lowers air and surface temperatures in densely populated regions.



Stormwater Reduction: Rainfall on impermeable surfaces, like concrete and asphalt, generates stormwater issues in cities, leading to problems such as flooding, water quality impairments, and reduced continuity of streamflow. In natural systems, rainwater interception and evapotranspiration minimize stormwater and reduce the reliance on costly engineered stormwater solutions.



Wildlife Habitat: Urban forests function as crucial wildlife habitats within the urban landscape, supporting a diverse range of species that have adapted to living alongside humans. These flora and fauna communities rely on these forests for essential resources, including refuge, food, water, and shelter, in an otherwise demanding environment.



Carbon Sequestration and Storage: Carbon dioxide (CO), the primary greenhouse gas driving global warming, is absorbed, and stored by trees during photosynthesis. This sequestered carbon is stored in the plant tissues during the lifetime of a tree.



Noise Buffering: Urban forests and tree canopies serve as natural noise buffers, reducing sound from traffic and other sources. The reduction of nuisance noise is beneficial to human health and well-being and can minimize noise impacts which negatively affect wildlife habitat.



Economic Benefits: Trees bring numerous economic advantages, such as higher property values, increased business traffic, heightened demand, tourism attraction, reduced energy costs, and resident appeal. Research indicates that urban forest programs typically yield substantial returns on investment, believed to be 2:1 or more (Endreny 2018).



Human Health and Wellness: Urban trees provide intangible yet significant societal benefits including recreation, enhancing the aesthetics of city streets, and fostering community pride and identity. Research also shows that trees play a role in improving health outcomes, reducing stress, enhancing mental well-being including cognition, attention, and anxiety, clinical outcomes, and crime reduction (Wolf et al. 2020).

Methods

i-Tree Study Design

The i-Tree Eco study was conducted using pre-stratified protocols to obtain representative samples with randomized 0.1-acre and 0.05-acre plots. Strata are consistent with the 2011 Lake Forest Park i-Tree study design for continuity in management units; these include parcels >¼-acres (large residential), parcels ≤⅓-acres (small residential), and the commercial town center. Road networks are excluded from sample selection since they are interwoven amongst other strata and incorporated into calculations for total strata area.

Plots are located on both public and private lands. To secure permission to collect data on private parcels, the City arborist, with support from DCG/Watershed staff, contacted landowners via mail, email, the City newsletter, and door-knocking. Additional randomly selected plots were generated in instances where permission was not granted, until the required number of research plots was reached.

Data from 100 plots were collected in 2022 and 2023. An additional 60 plots were planned in the study design but could not be collected due to being denied access onto private property.

Once processed with the user defined data and configuration, i-Tree provides statistical analysis and actionable insights on a range of urban forestry topics including structure and composition, benefits and costs, air quality interactions, and pest analysis. Analysis of invasive species was conducted using information from the King County Noxious Weed Board, and species designations recorded in the i-Tree Eco software were disregarded.

Stratum	Acres	Number of Plots
Town Center	19	8
≤¼ Acres	532	52
>¼ Acres	1750	40




Figure 2. Strata and plot location map.

i-Tree is a software suite and a set of tools developed by the USDA Forest Service and various partners to quantify the benefits and values of urban trees and forests. It provides a platform for assessing and managing urban forest ecosystems, focusing on the many environmental, economic, and societal benefits they offer.

i-Tree Manuals and Software Versions

- ❖i-Tree Software Suite v6.0
- ❖i-Tree Eco v6.0 User Manual
- i-Tree Eco v6.0 Field Manual




Urban Forest Measurements

This project utilizes data collection techniques as described in the i-Tree Eco v6.0 Field Manual. DCG/Watershed field researchers performed a range of measurements for each plot, encompassing both general plot characteristics and tree-specific measurement details. Plot-level and tree-level parameters are outlined below in the callout to the right. A total of 631 trees were assessed.

Limitations and Assumptions

Reported data was generated using i-Tree Eco, and therefore, limited by the associated model assumptions. Data provided by i-Tree Eco does not output standard error or other quantifiable metrics of sampling uncertainty for derived metrics. Standard error is reported for certain plot-level metrics supported by i-Tree Eco. Studies of i-Tree sampling methodology suggest that a 100-plot sample has an expected relative standard error (SE) of approximately 17%, however, this will differ by study and among assessed metrics (Nowak et al. 2008). Caution is advised in ascertaining trends between this study and the prior 2011 Lake Forest Park i-Tree Eco study for metrics which lack standard error metrics.



Plot Metrics

Plot ID

Date

Field Crew

Plot Center Address Coordinates (Lat/Long)

Tree Cover (%)

Shrub Cover (%)

Plantable Space (%)

Land Use

Ground Cover

Comments

Tree Metrics

Tree ID

Date

Status

Distance to Plot Center

Direction from Plot Center

Tree Species

DBH

Crown Condition (% Dieback)

Tree Height

Crown Top and Base Height

Crown Width (Bidirectional)

Percent of Crown Missing

Crown Light Exposure


Nearby Building Distance and

Direction

Street Tree

Comments

Section 9, ItemA.



Light Detection and Ranging (LiDAR) can be used to provide highly accurate and spatially explicit models of urban forests. Canopy height models (CHM) are useful as a tool in urban forest management to quantify forest structure. Pictured (left) is a graphic depicting a CHM model of tree canopy height. Other LiDAR applications in forestry include canopy cover analysis, forest health assessment, biomass and carbon estimation, tree inventory mapping, and urban planning and design.

Canopy Height Model

This assessment includes a canopy height model (CHM) analysis to provide information on urban forest structure and insight into retention of the City's largest and tallest trees. The CHM model utilized LiDAR data from the two most recent LiDAR flights on publicly available databases, 2016 and 2021.

Modeling was completed in the R Program using the 'lidR' package, an open-source software integrated into the R ecosystem, for the purpose of manipulating and visualizing LiDAR data with applications in forestry. Canopy height model and tree top identification algorithms were used to identify tree heights with a variable search window. Trees overlapping buildings were removed from the model output using Washington Department of Natural Resources Urban Forestry's 2022 King County Land Cover Metrics dataset and outliers below 15 feet in height were removed because could they not reliably distinguished from other shrubs or infrastructure.

This process yields a point layer with canopy height values that can be used to calculate derived metrics. This data is used to review trends of tree canopy height over time.



6

¹ LiDAR Data obtained from the Washington Department of Natural Resources LiDAR Portal. Sourced information includes 2016 data from Quantum Spatial and 2021 data from the Washington Geologic Survey.

Results

Tree Characteristics of the Urban Forest

Lake Forest Park is estimated to contain 297,056 (± 39,070 SE) trees, with canopy cover measuring 50.6% of the City area, ranking among the most heavily forested municipalities in the region. This is comparable to other canopy cover estimates including the study conducted by Elm in 2016, which estimated canopy cover of 50%, and i-Tree Landscape, which estimates canopy cover of 48% in 2017 (data obtained from i-Tree Landscape in November 2023). Similar results among differing methods validate the i-Tree Eco study findings.

The average tree density in Lake Forest Park is estimated to be 129 trees per acre (TPA). Large Residential areas have the highest tree density, followed by Small Residential areas, and Town Center.

Douglas-fir, bigleaf maple, and western red cedar continue to be the most common trees and are native to the Puget Lowlands Ecoregion. Diversity is key to resiliency in urban forests, particularly regarding impacts from disease and insects, and climate change.




Figure 3. Tree density by stratum.




Figure 4. Tree composition of common species.

Lake Forest Park has a greater canopy cover and tree density than any of the cities which i-Tree listed as comparable. Of these, Atlanta is reported to have the greatest tree canopy cover at 36.7% and Morgantown is reported to have the greatest tree density at 119 TPA.



Tree size class distributions provide a snapshot of forest structure that informs management strategies. Among these, it is useful to know whether a forest has a young or aging population. Currently, 71% of trees in Lake Forest Park are less than 12" diameter-at-breast-height (DBH) indicating a skew toward younger or smaller trees.

Despite a youthful population, or Type 1 distribution (Morgenroth et al. 2020), the tree size class distribution skews slightly larger than the prior 2011 i-Tree Eco study. The percentage of the largest trees, those above 30" DBH, have increased since 2011 and now account for 5% of trees. Trees greater than 24" inches DBH now account for 10% of the total tree population, an increase from 2011.

Trees in Lake Forest Park are estimated to be 63% native to Washington overall, concentrated most highly in the Large Residential stratum, followed by the Small Residential and Town Center Stratum (Figure 5).




Figure 6. Tree DBH class distribution by stratum.




Figure 5. Native status of trees by stratum.

Trees designated by King County or Washington State as noxious weeds comprise less than 1% of the tree population. These are represented by only one species, common hawthorn. However, 19% of trees are species listed by King County as weeds of concern. These include cherry laurel, bird cherry, European mountain ash, black locust, horse chestnut, and English holly.




Figure 7. Tree DBH distribution.

Urban Forest Cover and Leaf Area

Leaf area density is greatest in the Large Residential stratum compared to other strata due to high tree density and the presence of larger trees. As a result, leaf area in the large residential stratum is 3 times greater than the Small Residential stratum, and 7 times greater than the Town Center stratum (Figure 8). A handful of species contribute most of the leaf area including Douglas-fir, western red cedar, and bigleaf maple (Table 1). The importance value of each species represents the sum of the percent cover of a specific species and the leaf area percentage. This indicates which species dominate the urban canopy structure but are not always the best species to plant. The leaf area is an informative metric because it directly correlates with many urban forest functions and benefits such as avoided stormwater runoff.

Table 1. Leaf area, importance value, and percent of population by tree

Species	% Population	% Leaf Area	Importance Value
Douglas-fir	16.3	35.5	51.8
Bigleaf maple	11.1	18.9	30.0
Western red cedar	9.1	15.5	24.6
	-		-
Cherry laurel	8.2	0.7	8.9
Red alder	2.2	5.8	8.0
English holly	6.2	1.3	7.5
Vine maple	5.2	1.9	7.1
Bitter cherry	6.3	0.7	6.9
European bird	4.9	0.1	5.0
Deodar cedar	0.3	3.9	4.3
Western hemlock	2.8	1.3	4.1
Hinoki cypress	3.1	0.5	3.7
Japanese maple	2.8	0.5	3.3
Western white pine	0.4	2.8	3.2
Giant Sequoia spp	0.3	2.6	2.9
Northern white	2.3	0.1	2.3
Black poplar	1.2	1.0	2.2
Portugal laurel	1.9	0.3	2.2
Sitka spruce	0.3	1.6	2.0
Pacific dogwood	1.6	0.3	1.8
Plum spp	1.3	0.1	1.4
Lodgepole pine	0.7	0.4	1.1
Blue spruce	0.4	0.7	1.1



Total leaf area is defined as the one-sided area of all leaves in the study area. This differs from canopy cover because individual leaves may overlap within and among trees.



The total plantable space in Lake Forest Park is estimated to be $22.4\% \pm 2.9$ (SE), which represents opportunities for additional tree planting. This is defined as the amount of land area with suitable soils that are not under existing tree canopies or other overhead or land use restrictions that would prohibit tree planting (e.g., developed park or playfield).

Tree benefits are informed by groundcover composition since they interact with ground-level natural processes. Rainwater interception, for example, reduces runoff before entering stormwater systems.

Groundcover composition is consistent with expectations for the land use types, with high intensity land uses having the most buildings and impervious surfaces, and the low intensity land uses having the most groundcover vegetation, duff/mulch, and bare soil. Impervious surfaces are highest in the Town Center (82%), followed by Small Residential areas (51%), then Large Residential areas (25%).

Low building cover in the Town Center is believed to be due to error resulting from a low sample size, since much of the site appears to be composed of buildings based on visual estimates.




Figure 8. Leaf area density by stratum.




Figure 9. Ground cover composition by stratum.



Air Pollution

Many urban areas have high levels of air pollution which negatively impacts the health of humans and ecosystems. Urban forests mitigate the effects of air pollution through several processes including the absorption and particulate matter filtration, air temperature cooling, and reducing the energy consumption of buildings. While trees also emit volatile organic compounds (VOCs) that contribute to the formation of ozone (O), studies show that high tree cover is correlated with a reduction in ozone formation (Nowak and Dwyer, 2000).

The Lake Forest Park urban forest canopy is estimated to remove 1,607 pounds of carbon monoxide (CO), 33,013 pounds of nitrogen dioxide (NO₂), 69,299 pounds of O₃, 93,657 pounds of particulate matter less than 10 microns and greater than 2.5 microns (PM10), 12,458 pounds of particulate matter less than 2.5 microns (PM2.5), and 2,704 pounds of sulfur dioxide (SO₂) annually. This removal has an associated value of \$2.55 million.

Air pollution removal varies temporally, as shown in Figure 10. Some pollutants such as NO₂ and O₃ are removed at greater levels during the summer growing season while PM2.5 and PM10 removal is greatest during the fall and winter. Since some types of air pollution removal correlates with leaf area, the distribution of evergreen and deciduous trees also influences the magnitude of temporal variation.




Figure 10. Estimated monthly pollution removal by the Lake Forest Park urban forest.




Figure 11. Estimated monetary value of air pollution removal annually.



Carbon Sequestration and Storage

Tree canopy cover in Lake Forest Park is not just a local issue. Global climate change is largely driven by carbon dioxide (CO₂) emissions, a compound which trees uptake and sequester during photosynthesis. Carbon is stored in tree leaves and woody tissues, and therefore, reduces the amount of atmospheric carbon otherwise contributing to climate change. Carbon will remain in a tree until it eventually decomposes, where it may either be released to the atmosphere, returned to soil, or absorbed by other organisms.

The Lake Forest Park urban forest is estimated to remove 2,672 tons of carbon annually. Areas with

8,000

8,000

4,000

2,000

Large Residential Small Residential Town Center (>1/4 Acre)

Carbon Sequestration Density (lb/yr/ac) CO₂ Equivalent (lb/yr/ac)

Figure 12. Carbon sequestration each year by stratum.

more tree cover provide the greatest levels of CO₂ sequestration, such as the Large Residential stratum, which provide 2 to 6-fold more than the other strata on a per-area basis (Figure 12). The estimated value of this benefit is \$456,000 per year.

Carbon storage is also valuable to quantify because a tree that decomposes will eventually release CO₂ back into the atmosphere. Trees in Lake Forest Park collectively store 97,300 tons of carbon, with an estimated value of \$16.6 million. Douglas-fir, bigleaf maple, western red cedar, and black cottonwood are the tree species which currently have the greatest amount of carbon storage.




Figure 13. Total carbon storage by stratum.

Climate change is the process of shifting global and regional climate patterns, driven primarily by anthropogenic activities such as fossil fuel emissions and deforestation. These result in increased concentrations of greenhouse gases in the atmosphere which lead to globally rising temperatures, altered weather patterns, and sea level rise, which affect societies, economies, and ecosystems across the planet. Changing climates also mean cities need to manage for resilient forests which can tolerate shifting conditions.

Surface Water Runoff

Runoff from impermeable surfaces is a significant source of water pollution and flooding, posing risks to both human and environmental well-being while imposing substantial economic costs. Trees play a role in mitigating runoff through evapotranspiration, a combination of processes which include the interception of rainwater, evaporation, and transpiration, and thereby, return water to the atmosphere. Additionally, trees enhance the ability of rainwater to infiltrate into soils through inputs of organic matter and improving porosity. The combination of these processes results in the attenuation of pollution laden runoff and reduction in the severity of flooding events. Urban forests in Lake Forest Park are estimated to reduce runoff by 50.4 million gallons per year.

Urban forests also reduce the need for cities to rely on costly built infrastructure to manage water quality and quantity issues. This "green infrastructure" is estimated to provide Lake Forest Park with estimated economic benefit at \$450,000 per year for water quality and flood reduction benefits they provide. The majority of these benefits are provided in the Large Residential stratum, where tree density and leaf area are greatest.




Figure 14. Avoided runoff per annum, by stratum.






Figure 15. Value of avoided runoff per annum, by stratum.

Tree Benefits Summary

The total economic benefit of trees in Lake Forest Park is estimated to be \$4.1 million per year, when accounting for energy savings, gross carbon sequestration, pollution removal, and avoided runoff (Table 2). On an individual basis, this amounts to \$13.79 per tree.

Tree replacement values are another useful measure when managing forests since it is more expensive to replace trees than preserve existing trees. The collective replacement value of all trees in Lake Forest Park is estimated to be \$531 million in addition to the \$16.9 million provided by carbon storage. The high cost of tree removal can inform public policy and management decisions regarding tree preservation and replacement on public and private land.

Trees in Lake Forest Park also generate 5,402 tons of oxygen every year, however, this benefit is believed to be relatively insignificant due to the vast reserves of oxygen in the atmosphere and production from oceanic systems (Broecker, 1970; i-Tree, 2023).

Table 2. Total Lake Forest Park tree benefits summary.

Benefits	Annual Value	Annual Value Per Tree
Energy & Carbon Emission Reduction	\$646,000	\$2.17
Gross Carbon Sequestration	\$455,648	\$1.53
Pollution Removal	\$2,545,701	\$8.57
Avoided Runoff	\$450,254	\$1.52
Total Benefits	\$4,097,603	\$13.79

Urban forests result in a net reduction in energy use through shading, evaporative cooling, and blocking of winter winds which are estimated to save Lake Forest Park residents \$542,683 per year. Additionally, the value of reduced carbon emissions resulting from energy savings is valued at \$104,000 per year.





Pests and Pathogens

Trees are susceptible to pests and pathogens that are capable of impacting tree viability, resulting in reduced lifespan, hazard conditions and sometimes mortality. The i-Tree Eco model included an analysis of the of Lake susceptibility Forest Park's urban forests common pests and pathogens to evaluate risks and management priorities. Of these, 15 currently present in King County based on the pest range maps developed by the Forest Health Technology Enterprise Team (i-

Tree 2023), as shown in Figure 16. See Appendix V for the complete list of pests and pathogens assessed through i-Tree. Of the 15 pests present in King County, we have highlighted the top three that could impact the dominant canopy species within Lake Forest Park. This includes two fungal pathogens and one insect commonly found in Pacific Northwest forests, Armillaria root disease, Heterobasidion root disease, and western spruce




Figure 16. Susceptibility by trees to the 15 evaluated pests and pathogens which are currently known to be present in King County.

budworm. It's important to note that some pests and pathogens are naturally occurring and play an important role in forest ecological processes, while others have significant negative ecological and economic impacts.

Armillaria Root Disease (*Armillaria sp.*) refers to a group of fungi that causes reduced leader growth and foliage discoloration and thinning, spreading

Pests and pathogens included in the i-Tree analysis have been documented within King County limits but does not confirm their presence in the trees surveyed within the study plots. This research did not include an advanced level of tree health analysis beyond the standard i-Tree data collection protocols.

through a tree's root system (Allen et al. 1996). Trees susceptible to Armillaria documented in the Lake Forest Park study include Douglas fir, subalpine fir, western red cedar, but can also impact broadleaved trees as well. This represents 25.5% of Lake Forest Park's urban forest and \$11.4 million in replacement value (i-Tree 2023).

Heterobasidion Root Disease (HRD; Heterobasidion annosum, H. occidentale) - also called Annosus root and butt rot – is a fungus known to impact many of our native conifers as well as bigleaf maple and alder. In younger trees, symptoms include a reduction in the leader and branch growth, chlorotic foliage and a distressed cone crop. Trees become infected by airborne spores that may enter through wounds on branches, trunks or roots, but can then spread from tree to tree via root systems (Allen et al. 1996). Trees present in Lake Forest Park which are susceptible to HRD include Douglas-fir, western white pine, subalpine fir, Norway spruce, western hemlock, shore pine, and western red cedar. This represents 29.4% of the City's trees with a replacement value of \$17.95 million (i-Tree 2023).



Photo by USDA Forest Service; fruiting body of H. occidentale.

Western spruce budworm (Choristoneura occidentalis, C. freemani) is an insect native to western North America and is a widespread defoliator of several native conifer species. It feeds upon and

defoliates Douglas-firs, spruce, and the true firs (e.g., white fir, subalpine fir). The larvae feed on the current year's needles and buds giving the canopy a red-brown or grayish appearance with thinning foliage and produces a new generation annually. It is typically controlled through natural predators but can be controlled through insecticides (Fellin and Dewey 1986). Per this i-Tree survey, 20.1% of the tree population is susceptible and has a replacement value of \$13.5 million (i-Tree 2023).



Photo by: Montana State University Extension; Western spruce budworm.

Emerging Threats in Western Washington

Disease and pest outbreaks have increased in number and frequency in recent years due to international trade, travel, and climatic changes. New pests are introduced outside of their native range into ecosystems that have not evolved with the pest to develop any resistance. Climate changes, such as increases in seasonal and average air temperatures, increases in extreme heat, and prolonged drought, add abiotic stressors—weakening a tree's ability to defend against these diseases and pest pressures (Mauger et al. 2015). The Pacific Northwest region currently faces several emerging threats, namely sooty bark disease, bronze

birch borer, emerald ash borer, and non-native longhorned beetle species.

Sooty bark disease (Cryptostroma corticale) causes dieback primarily in maple species. To date, the fungus has been found to cause damage in sycamore maples (Acer pseudoplatanus), red maple (A. rubrum), Japanese maple (A. palmatum), vine maple (A. circinatum), and bigleaf maple (A. macrophyllum) in the Puget Sound region. Other confirmed hosts include Pacific dogwood (Cornus nuttallii) and horse chestnut (Aesculus hippocastanum) (Brooks et al. 2022). The fungus infects the tree's vascular system and thrives during hot summers, proliferating in drought-stressed trees (Brooks et al. 2022). Sooty bark disease was not included in the i-Tree replacement value analysis.

Bronze birch borer (*Agrilus anxius*) is a beetle whose larvae tunnel into live wood, creating extensive galleries leading to branch or trunk girdling, ultimately cutting the rest of the branch off from resources. Bronze birch borers are attracted to trees weakened by environmental stressors, age, or other diseases and pests (Antonelli 2008). Paper birch (*Betula papyrifera*), European white birch (*B. pendula*), and grey birch (*B. populifolia*) are more susceptible than other birch species. Bronze birch borer was not included in the i-Tree replacement value analysis.

Emerald ash borer (*Agrilus planipennis*) has been present in the United States since 2002 but only recently has been confirmed in the Pacific Northwest Region since 2022, where it was discovered in Oregon. While it has not yet been sighted in the Puget Sound region, its spread into Washington State is expected. The emerald ash borer infects native and non-native ash trees (*Fraxinus* spp.). The beetles do not discern between stressed or healthy trees, and the impact is anticipated to be significant, especially in native forests. Like other borers, its larvae create extensive


galleries, causing limb and trunk dieback leading to decline and eventual tree death (Bliss-Ketchum et al. 2021). Although no ash trees were identified in this study, they are likely present in the City. Oregon ash (*Fraxinus latifolia*) is a native tree which can be found near water or in wetland areas. Ashes are also commonly planted as street trees and as ornamentals in yards and gardens.



Photo by Leah Bauer, USDA Forest Service, Northern Research Center Station; Emerald ash borer

Asian, citrus, and red-necked long-horned beetles

(Anoplophora glabripennis, A. chinenses, and Aromia bungii respectively) feed on the wood of hardwood trees. Although there are no known established populations of these beetles in Washington, they have reached local nurseries where they were eradicated. With continued global movement within the nursery trade, Washington will need to continually monitor these species. The beetles typically feed on both healthy and dying trees and are known to impact 40 host species including maples, horse chestnuts, willows, birches, and elms. There are locally known native look-a-likes which present challenge identification for to nonprofessionals (WISC 2017a). The Washington Invasive Species Council has resources for identifying the potentially invasive versus native beetles in King County. Asian long-horned beetle is expected to impact 66,215 trees in Lake Forest Park with a replacement value of \$1.91 million (i-Tree 2023).



Canopy Height Model Results

The distribution of tree heights in Lake Forest Park reveals that the proportion of tall trees, those greater than 135 feet in height, have increased by 21% in 2021 compared to 2016. The proportion of the tallest trees, those greater than 165 feet increased by 86% during this period, albeit accounting for less than 1% of the total tree population. This suggests that most tall trees are being retained, and that other small and moderate size trees are aging into the larger height classifications. The tallest tree is estimated to be 195 feet tall.

Trends of smaller trees vary by height class, although the proportion of trees in the moderate

height classes have tended to decrease while the 15-30 smallest, those between feet, are approximately equal. Since trees below 15 feet were removed in this analysis, plot samples collected as part of the i-Tree Eco inventory provide better insight into age distribution and forest regeneration. The canopy height model is less selective than the plot sampling method in finding smaller trees and subcanopy trees, so interpretations of age and regeneration are not as precise as other sampling methods. However, this analysis provides us with additional insight into the distribution of trees within the assessed range between 15 and 195 feet.




Figure 17. Histogram of tree heights using LiDAR data from 2016 and 2021.



Discussion

The results of this i-Tree Eco study and canopy height model provide insight on the current composition and structure of Lake Forest Park's urban and community forest as well as quantify ecosystem service benefits and values. The results suggest a net increase in urban tree canopy cover and tree density during the last ten years and an increasing trend in the presence of large canopy trees, primarily comprised of Pacific Northwest native species.

The data provided by this study provides City urban forest managers with practical information that is useful to develop urban forest management strategies and policies. Cities across the Puget Sound region and the Pacific Northwest face several challenges to steward resilient, regenerative, and viable urban forests. These include shifts in climate conditions, threats from current and emerging pests and pathogens, the potential for increases in urban wildfires, and continued development needed to meet regional housing needs. Urban forest managers are also tasked with ensuring that tree canopy remains equitably distributed throughout the City and that more densely developed land use zones have adequate green infrastructure to manage stormwater, minimize urban heat islands, provide shade, and foster both ecological health as well as human health and wellness.

Climate Adaptation and Resilience

Within the field of urban forest management, arborists, ecologists, foresters, and land managers continue to evaluate best management practices and adapt arboricultural strategies to the on-the-ground conditions impacting the resilience of urban forests.

Western Washington is expected to experience drier conditions increasingly and higher temperatures during the summer months, with potential increases in precipitation during the winter months. This will present and exacerbate stressors on existing urban forests such as drought, insect and tree disease outbreaks, competition with invasive plant species, habitat loss and fragmentation, erosion, and wildfires. This also creates challenges for establishing the next generation of urban forest canopy, especially coupled with development pressures and the need to respond to the rising need for sustainable and affordable housing.

One strategy for establishing resilience within the urban forest is to increase species diversity (at the family, genus, and species level), ensure installed trees are climate adapted to current and future stressors, such as drought. Since most biotic and abiotic stressors exhibit variable effects among tree species, a diverse forest acts as an insurance policy that minimizes risk from impacts to individual taxa. The City of Lake Forest Park currently has an approved tree list which includes species that are "better performing" in the built environment and drought tolerant. This is an important educational and management tool that should be periodically evaluated and updated to account for updated research and recommendations from the arboriculture and horticultural trades and account for climate resilience.

Protection of Significant and Large Diameter Trees

The tree size class distributions outlined in this study tell us that 71% of the City's forests are less than 12" diameter-at-breast-height (DBH), but that the percentage of large diameter trees (those greater than 24"DBH) has continued to increase during the last decade. Tree diameter correlates with tree height and volume and can be used as a metric to describe overall tree size and identify large trees, which are a management priority for the City. Large trees provide greater levels of ecosystem services such as stormwater capture and infiltration, cooling, and water quality improvements compared to small trees; and therefore, societal benefits are optimized when they are retained.

Since the majority of Lake Forest Park's urban forest is located on private residential, commercial, and industrial property, protection of significant and large diameter trees on privately owned property



will be an important strategy as the City seeks to protect its existing tree canopy. The City currently regulates trees during development of private property through its tree ordinance – Chapter 16.14 Tree Canopy Preservation and Enhancement - as well as trees within shoreline jurisdiction through Chapter 16.18 Shoreline Master Program. These regulations prioritize the retention and protection of existing trees and groves as well as replanting with new trees when removal is unavoidable due to tree risks, site development design, and storm damage.

The findings within this report can be used as a tool to educate and engage community members, private landowners, and the development community to encourage early assessment and integration of existing significant and exceptional trees in the predesign or early design phase of new development. Another critical component is ensuring not only the long-term viability of a retained tree but ensuring that replacement trees are chosen using the "Right Plan, Right Place" approach, have adequate growing conditions (e.g., soil volumes, planter strip widths etc.) to reach maturity without impacting required infrastructure such as sidewalks, driveways, and utilities.

Invasive Species Management

Of the trees surveyed, 19% are either listed as a Class C noxious weed (common hawthorn) or a Weed of Concern (bird cherry, black locust, cherry laurel, English holly, and European mountain ash) by the King County Noxious Weed Program. These species are "non-regulated" meaning that property owners in King County are not required to control these species, but control is recommended where feasible. Cherry laurel and English holly, cherry laurel, black locust, and European mountain ash are widely used as ornamental landscape plants. However, these species compete with our native

flora and naturalize in open spaces and critical areas.

A potential strategy to address this problem could be for the City to develop a prohibited species list and other educational materials to discourage property owners and developers from introducing these species into new plantings. In addition, the City could consider removal of these species from public open spaces, replacing them with native tree species.

Additional Considerations

Scientific studies as well as programmatic and policy audits provide important data to evaluate the success of urban forest resource management strategies. In addition to repeating this study, evaluation of other policies and regulations can inform municipal code updates, the effectiveness of current community education and outreach efforts, and additional support needed from community members in managing trees on their properties. Continued study of on-the-ground conditions coupled with evaluation of existing policies and best practices will provide the City with the tools and information needed to effectively and adaptively manage the valuable urban forest resource.



References

- Allen, E.A., D.J. Morrison, and G.W. Wallis. (1996). *Common Tree Diseases of British Columbia*. Natural Resources Canada/Canadian Forest Service.
- Antonelli, A. (2008). *Bronze Birch Borer* (Report No. EB1380E). Washington State University Extension. EB1380E. https://pubs.extension.wsu.edu/bronze-birch-borer
- Brooks, B., Hulbert, J.M., Omdal, D., Elliot, M., and Chastagner, G. A. (2022). *Sooty bark disease diagnostic guide* (Report No. FS375E). Washington State University Extension. https://pubs.extension.wsu.edu/sooty-bark-disease-diagnostic-guide
- Bliss-Ketchum L.B., R. Draheim, M. Hepner, and O. Guethling. (2021). *Emerald Ash Borer: Readiness and Response Plan for Oregon*. EAB Readiness and Response Steering Committee. https://www.oregon.gov/odf/Documents/forestbenefits/eab-readiness-and-response-plan-for-oregon.pdf.
- Broecker, W. S. (1970). A boundary condition on the evolution of atmospheric oxygen. *Journal of Geophysical Research*, 75(18), 3553–3557. https://doi.org/10.1029/JC075i018p03553
- Elm. (2016). Tree Canopy Extent, City of Lake Forest Park Tree Canopy Study.
- Endreny, T. A. (2018). Strategically growing the urban forest will improve our world. *Nature Communications*, 9(1), 1160–1163. https://doi.org/10.1038/s41467-018-03622-0
- City of Lake Forest Park. (2010). Community Forest Management Plan. https://www.cityoflfp.gov/DocumentCenter/View/6175/Community-Forest-Management-Plan?bidId=
- City of Lake Forest Park. (2011). *I-Tree Ecosystem Analysis: Urban Forest Effects and Values*. i-Tree Report. https://www.cityoflfp.gov/DocumentCenter/View/385/itreereport?bidId=
- City of Lake Forest Park. (n.d.). *General tree list for lake forest park, WA*. Accessed November 2023. https://www.cityoflfp.gov/DocumentCenter/View/224/Tree-List?bidId=
- City of Lake Forest Park. (2023a). *Municipal Code Chapter 16.14 Tree Canopy Preservation and Enhancement*. Accessed October 2023. https://www.codepublishing.com/WA/LakeForestPark/
- City of Lake Forest Park. (2023b). *Municipal Code Chapter 16.18 Shoreline Master Program*. Accessed October 2023. https://www.codepublishing.com/WA/LakeForestPark/
- i-Tree. (2021a). Field Guide. https://www.itreetools.org/documents/274/EcoV6.FieldManual.2021.10.06.pdf
- i-Tree. (2021b). *User's Manual*. https://www.itreetools.org/documents/275/EcoV6_UsersManual.2021.09.22.pdf
- i-Tree. (2023). i-Tree Eco Software Suite v6.0.
- King County Noxious Weed Control Board. (2023). *King County Noxious Weed List*. https://your.kingcounty.gov/dnrp/library/water-and-land/weeds/weedlists/2023-Official-Weed-List_common-name.pdf

- Mauger, G.S., J.H. Casola, H.A. Morgan, R.L. Strauch, B. Jones, B. Curry, T.M. Busch Isaksen, L. Whitely Binder, M.B. Krosby, and A.K. Snover, 2015. *State of Knowledge: Climate Change in Puget Sound Report* prepared for the Puget Sound Partnership and the National Oceanic and Atmospheric Administration. Climate Impacts Group, University of Washington.
- Morgenroth, J., Nowak, D. J., & Koeser, A. K. (2020). DBH Distributions in America's Urban Forests-An Overview of Structural Diversity. *Forests*, 11(2), 135-. https://doi.org/10.3390/f11020135
- Nowak, D. J., & Dwyer, J. F. (2000). *Understanding the Benefits and Costs of Urban Forest Ecosystems*. In Handbook of Urban and Community Forestry in the Northeast (pp. 11–25). Springer US. https://doi.org/10.1007/978-1-4615-4191-2_2
- Nowak, D., Walton, J., Stevens, J., Crane, D., & Hoehn, R. (2008). Effect of Plot and Sample Size on Timing and Precision of Urban Forest Assessments. *Arboriculture & Urban Forestry*, 34(6), 386–390. https://doi.org/10.48044/jauf.2008.052
- Fellin, D.G., and Dewey, J.E. (1986). *Western Spruce Budworm* (Forest Insect and Diseases Leaflet 53). U.S. Forest Service. https://www.fs.usda.gov/Internet/FSE_DOCUMENTS/fsbdev2_043445.pdf
- Washington Invasive Species Council (WISC). (2017a). *Citrus, Asian, and Red-Necked Longhorned Beetles Fact Sheet*. https://invasivespecies.wa.gov/wp-content/uploads/2019/07/InvasiveLonghornedBeetles-FactSheet.pdf
- Washington Invasive Species Council (WISC). (2017b). *Emerald Ash Borer, Stop the Invasion*. https://invasivespecies.wa.gov/wp-content/uploads/2019/07/EmeraldAshBorer-FactSheet.pdf
- Wolf, K. L., Lam, S. T., McKeen, J. K., Richardson, G. R. A., van den Bosch, M., & Bardekjian, A. C. (2020). Urban Trees and Human Health: A Scoping Review. *International Journal of Environmental Research and Public Health*, 17(12), 4371-. https://doi.org/10.3390/ijerph17124371

The following appendices are provided by the i-Tree Eco software as a component of the madel outp

Section 9. ItemA.

Appendix I. i-Tree Eco Model and Field Measurements

i-Tree Eco is designed to use standardized field data from randomly located plots and local hourly air pollution and meteorological data to quantify urban forest structure and its numerous effects (Nowak and Crane 2000), including:

- Urban forest structure (e.g., species composition, tree health, leaf area, etc.).
- Amount of pollution removed hourly by the urban forest, and its associated percent air quality improvement throughout a year.
- Total carbon stored and net carbon annually sequestered by the urban forest.
- Effects of trees on building energy use and consequent effects on carbon dioxide emissions from power sources.
- Replacement value of the forest, as well as the value for air pollution removal and carbon storage and sequestration.
- Potential impact of infestations by pests, such as Asian longhorned beetle, emerald ash borer, gypsy moth, and Dutch elm disease.

Typically, all field data are collected during the leaf-on season to properly assess tree canopies. Typical data collection (actual data collection may vary depending upon the user) includes land use, ground and tree cover, individual tree attributes of species, stem diameter, height, crown width, crown canopy missing and dieback, and distance and direction to residential buildings (Nowak et al 2005; Nowak et al 2008).

During data collection, trees are identified to the most specific taxonomic classification possible. Trees that are not classified to the species level may be classified by genus (e.g., ash) or species groups (e.g., hardwood). In this report, tree species, genera, or species groups are collectively referred to as tree species.

Tree Characteristics:

Leaf area of trees was assessed using measurements of crown dimensions and percentage of crown canopy missing. In the event that these data variables were not collected, they are estimated by the model.

An analysis of invasive species is not available for studies outside of the United States. For the U.S., invasive species are identified using an invasive species list (Oregon Invasive Species Council 2014) for the state in which the urban forest is located. These lists are not exhaustive and they cover invasive species of varying degrees of invasiveness and distribution. In instances where a state did not have an invasive species list, a list was created based on the lists of the adjacent states. Tree species that are identified as invasive by the state invasive species list are cross-referenced with native range data. This helps eliminate species that are on the state invasive species list, but are native to the study area.

Air Pollution Removal:

Pollution removal is calculated for ozone, sulfur dioxide, nitrogen dioxide, carbon monoxide, particulate matter less than 2.5 microns, and particulate matter less than 10 microns and greater than 2.5 microns. PM2.5 is generally more relevant in discussions concerning air pollution effects on human health.

Air pollution removal estimates are derived from calculated hourly tree-canopy resistances for ozone, and sulfur and nitrogen dioxides based on a hybrid of big-leaf and multi-layer canopy deposition models (Baldocchi 1988; Baldocchi et al 1987). As the removal of carbon monoxide and particulate matter by vegetation is not directly related to transpiration, removal rates (deposition velocities) for these pollutants were based on average measured values from the literature (Bidwell and Fraser 1972; Lovett 1994) that were adjusted depending on leaf phenology and leaf area. Particulate removal incorporated a 50 percent resuspension rate of particles back to the atmosphere (Zinke 1967). Recent updates (2011) to air quality modeling are based on improved leaf area index simulations, weather and

Section 9, ItemA.

pollution processing and interpolation, and updated pollutant monetary values (Hirabayashi et al 201 et al 2012; Hirabayashi 2011).

Trees remove PM2.5 and PM10* when particulate matter is deposited on leaf surfaces (Nowak et al 2013). This deposited PM2.5 and PM10* can be resuspended to the atmosphere or removed during rain events and dissolved or transferred to the soil. This combination of events can lead to positive or negative pollution removal and value depending on various atmospheric factors. Generally, PM2.5 and PM10* removal is positive with positive benefits. However, there are some cases when net removal is negative or resuspended particles lead to increased pollution concentrations and negative values. During some months (e.g., with no rain), trees resuspend more particles than they remove. Resuspension can also lead to increased overall PM2.5 and PM10* concentrations if the boundary layer conditions are lower during net resuspension periods than during net removal periods. Since the pollution removal value is based on the change in pollution concentration, it is possible to have situations when trees remove PM2.5 and PM10* but increase concentrations and thus have negative values during periods of positive overall removal. These events are not common, but can happen.

For reports in the United States, default air pollution removal value is calculated based on local incidence of adverse health effects and national median externality costs. The number of adverse health effects and associated economic value is calculated for ozone, sulfur dioxide, nitrogen dioxide, and particulate matter less than 2.5 microns using data from the U.S. Environmental Protection Agency's Environmental Benefits Mapping and Analysis Program (BenMAP) (Nowak et al 2014). The model uses a damage-function approach that is based on the local change in pollution concentration and population. National median externality costs were used to calculate the value of carbon monoxide removal (Murray et al 1994).

For international reports, user-defined local pollution values are used. For international reports that do not have local values, estimates are based on either European median externality values (van Essen et al 2011) or BenMAP regression equations (Nowak et al 2014) that incorporate user-defined population estimates. Values are then converted to local currency with user-defined exchange rates.

For this analysis, pollution removal value is calculated based on the prices of \$1,397 per ton (carbon monoxide), \$4,926 per ton (ozone), \$613 per ton (nitrogen dioxide), \$181 per ton (sulfur dioxide), \$330,079 per ton (particulate matter less than 2.5 microns), \$6,565 per ton (particulate matter less than 10 microns and greater than 2.5 microns).

Carbon Storage and Sequestration:

Carbon storage is the amount of carbon bound up in the above-ground and below-ground parts of woody vegetation. To calculate current carbon storage, biomass for each tree was calculated using equations from the literature and measured tree data. Open-grown, maintained trees tend to have less biomass than predicted by forest-derived biomass equations (Nowak 1994). To adjust for this difference, biomass results for open-grown urban trees were multiplied by 0.8. No adjustment was made for trees found in natural stand conditions. Tree dry-weight biomass was converted to stored carbon by multiplying by 0.5.

Carbon sequestration is the removal of carbon dioxide from the air by plants. To estimate the gross amount of carbon sequestered annually, average diameter growth from the appropriate genera and diameter class and tree condition was added to the existing tree diameter (year x) to estimate tree diameter and carbon storage in year x+1.

Carbon storage and carbon sequestration values are based on estimated or customized local carbon values. For international reports that do not have local values, estimates are based on the carbon value for the United States (U.S. Environmental Protection Agency 2015, Interagency Working Group on Social Cost of Carbon 2015) and converted to local currency with user-defined exchange rates.

For this analysis, carbon storage and carbon sequestration values are calculated based on \$171 per ton.

Oxygen Production:

The amount of oxygen produced is estimated from carbon sequestration based on atomic weights: net O2 release (kg/yr) = net C sequestration $(kg/yr) \times 32/12$. To estimate the net carbon sequestration rate, the amount of carbon sequestered as a result of tree growth is reduced by the amount lost resulting from tree mortality. Thus, net carbon sequestration and net annual oxygen production of the urban forest account for decomposition (Nowak et al 2007). For complete inventory projects, oxygen production is estimated from gross carbon sequestration and does not account for decomposition.

Avoided Runoff:

Annual avoided surface runoff is calculated based on rainfall interception by vegetation, specifically the difference between annual runoff with and without vegetation. Although tree leaves, branches, and bark may intercept precipitation and thus mitigate surface runoff, only the precipitation intercepted by leaves is accounted for in this analysis.

The value of avoided runoff is based on estimated or user-defined local values. For international reports that do not have local values, the national average value for the United States is utilized and converted to local currency with user-defined exchange rates. The U.S. value of avoided runoff is based on the U.S. Forest Service's Community Tree Guide Series (McPherson et al 1999; 2000; 2001; 2002; 2003; 2004; 2006a; 2006b; 2006c; 2007; 2010; Peper et al 2009; 2010; Vargas et al 2007a; 2007b; 2008).

For this analysis, avoided runoff value is calculated based on the price of \$0.01 per gallon.

Building Energy Use:

If appropriate field data were collected, seasonal effects of trees on residential building energy use were calculated based on procedures described in the literature (McPherson and Simpson 1999) using distance and direction of trees from residential structures, tree height and tree condition data. To calculate the monetary value of energy savings, local or custom prices per MWH or MBTU are utilized.

For this analysis, energy saving value is calculated based on the prices of \$96.70 per MWH and \$10.65 per MBTU.

Replacement Values:

Replacement value is the value of a tree based on the physical resource itself (e.g., the cost of having to replace a tree with a similar tree). Replacement values were based on valuation procedures of the Council of Tree and Landscape Appraisers, which uses tree species, diameter, condition, and location information (Nowak et al 2002a; 2002b). Replacement value may not be included for international projects if there is insufficient local data to complete the valuation procedures.

Potential Pest Impacts:

The complete potential pest risk analysis is not available for studies outside of the United States. The number of trees at risk to the pests analyzed is reported, though the list of pests is based on known insects and disease in the United States.

For the U.S., potential pest risk is based on pest range maps and the known pest host species that are likely to experience mortality. Pest range maps for 2012 from the Forest Health Technology Enterprise Team (FHTET) (Forest Health Technology Enterprise Team 2014) were used to determine the proximity of each pest to the county in which

Section 9, ItemA.

within 250 miles of the county edge, is between 250 and 750 miles away, or is greater than 750 miles away. FHTET did not have pest range maps for Dutch elm disease and chestnut blight. The range of these pests was based on known occurrence and the host range, respectively (Eastern Forest Environmental Threat Assessment Center; Worrall 2007).

Relative Tree Effects:

The relative value of tree benefits reported in Appendix II is calculated to show what carbon storage and sequestration, and air pollutant removal equate to in amounts of municipal carbon emissions, passenger automobile emissions, and house emissions.

Municipal carbon emissions are based on 2010 U.S. per capita carbon emissions (Carbon Dioxide Information Analysis Center 2010). Per capita emissions were multiplied by city population to estimate total city carbon emissions.

Light duty vehicle emission rates (g/mi) for CO, NOx, VOCs, PM10, SO2 for 2010 (Bureau of Transportation Statistics 2010; Heirigs et al 2004), PM2.5 for 2011-2015 (California Air Resources Board 2013), and CO2 for 2011 (U.S. Environmental Protection Agency 2010) were multiplied by average miles driven per vehicle in 2011 (Federal Highway Administration 2013) to determine average emissions per vehicle.

Household emissions are based on average electricity kWh usage, natural gas Btu usage, fuel oil Btu usage, kerosene Btu usage, LPG Btu usage, and wood Btu usage per household in 2009 (Energy Information Administration 2013; Energy Information Administration 2014)

- CO2, SO2, and NOx power plant emission per KWh are from Leonardo Academy 2011. CO emission per kWh
 assumes 1/3 of one percent of C emissions is CO based on Energy Information Administration 1994. PM10
 emission per kWh from Layton 2004.
- CO2, NOx, SO2, and CO emission per Btu for natural gas, propane and butane (average used to represent LPG), Fuel #4 and #6 (average used to represent fuel oil and kerosene) from Leonardo Academy 2011.
- CO2 emissions per Btu of wood from Energy Information Administration 2014.
- CO, NOx and SOx emission per Btu based on total emissions and wood burning (tons) from (British Columbia Ministry 2005; Georgia Forestry Commission 2009).

Appendix II. Relative Tree Effects

The urban forest in Lake Forest Park Plot Inventory 2023 provides benefits that include carbon storage and sequestration, and air pollutant removal. To estimate the relative value of these benefits, tree benefits were compared to estimates of average municipal carbon emissions, average passenger automobile emissions, and average household emissions. See Appendix I for methodology.

Carbon storage is equivalent to:

- Amount of carbon emitted in Lake Forest Park Plot Inventory 2023 in 492 days
- Annual carbon (C) emissions from 68,800 automobiles
- Annual C emissions from 28,200 single-family houses

Carbon monoxide removal is equivalent to:

- Annual carbon monoxide emissions from 7 automobiles
- Annual carbon monoxide emissions from 20 single-family houses

Nitrogen dioxide removal is equivalent to:

- · Annual nitrogen dioxide emissions from 2,360 automobiles
- Annual nitrogen dioxide emissions from 1,060 single-family houses

Sulfur dioxide removal is equivalent to:

- Annual sulfur dioxide emissions from 14,500 automobiles
- Annual sulfur dioxide emissions from 38 single-family houses

Annual carbon sequestration is equivalent to:

- Amount of carbon emitted in Lake Forest Park Plot Inventory 2023 in 14.0 days
- Annual C emissions from 1,900 automobiles
- Annual C emissions from 800 single-family houses

Appendix III. Comparison of Urban Forests

A common question asked is, "How does this city compare to other cities?" Although comparison among cities should be made with caution as there are many attributes of a city that affect urban forest structure and functions, summary data are provided from other cities analyzed using the i-Tree Eco model.

I. City totals for trees

City	% Tree Cover	Number of Trees	Carbon Storage	Carbon Sequestration	Pollution Removal
			(tons)	(tons/yr)	(tons/yr)
Toronto, ON, Canada	26.6	10,220,000	1,221,000	51,500	2,099
Atlanta, GA	36.7	9,415,000	1,344,000	46,400	1,663
Los Angeles, CA	11.1	5,993,000	1,269,000	77,000	1,975
New York, NY	20.9	5,212,000	1,350,000	42,300	1,676
London, ON, Canada	24.7	4,376,000	396,000	13,700	408
Chicago, IL	17.2	3,585,000	716,000	25,200	888
Phoenix, AZ	9.0	3,166,000	315,000	32,800	563
Baltimore, MD	21.0	2,479,000	570,000	18,400	430
Philadelphia, PA	15.7	2,113,000	530,000	16,100	575
Washington, DC	28.6	1,928,000	525,000	16,200	418
Oakville, ON , Canada	29.1	1,908,000	147,000	6,600	190
Albuquerque, NM	14.3	1,846,000	332,000	10,600	248
Boston, MA	22.3	1,183,000	319,000	10,500	283
Syracuse, NY	26.9	1,088,000	183,000	5,900	109
Woodbridge, NJ	29.5	986,000	160,000	5,600	210
Minneapolis, MN	26.4	979,000	250,000	8,900	305
San Francisco, CA	11.9	668,000	194,000	5,100	141
Morgantown, WV	35.5	658,000	93,000	2,900	72
Moorestown, NJ	28.0	583,000	117,000	3,800	118
Hartford, CT	25.9	568,000	143,000	4,300	58
Jersey City, NJ	11.5	136,000	21,000	890	41
Casper, WY	8.9	123,000	37,000	1,200	37
Freehold, NJ	34.4	48,000	20,000	540	22

II. Totals per acre of land area

City	Number of Trees/ac	Carbon Storage	Carbon Sequestration	Pollution Removal
		(tons/ac)	(tons/ac/yr)	(lb/ac/yr)
Toronto, ON, Canada	64.9	7.8	0.33	26.7
Atlanta, GA	111.6	15.9	0.55	39.4
Los Angeles, CA	19.6	4.2	0.16	13.1
New York, NY	26.4	6.8	0.21	17.0
London, ON, Canada	75.1	6.8	0.24	14.0
Chicago, IL	24.2	4.8	0.17	12.0
Phoenix, AZ	12.9	1.3	0.13	4.6
Baltimore, MD	48.0	11.1	0.36	16.6
Philadelphia, PA	25.1	6.3	0.19	13.6
Washington, DC	49.0	13.3	0.41	21.2
Oakville, ON , Canada	78.1	6.0	0.27	11.0
Albuquerque, NM	21.8	3.9	0.12	5.9
Boston, MA	33.5	9.1	0.30	16.1
Syracuse, NY	67.7	10.3	0.34	13.6
Woodbridge, NJ	66.5	10.8	0.38	28.4
Minneapolis, MN	26.2	6.7	0.24	16.3
San Francisco, CA	22.5	6.6	0.17	9.5
Morgantown, WV	119.2	16.8	0.52	26.0
Moorestown, NJ	62.1	12.4	0.40	25.1
Hartford, CT	50.4	12.7	0.38	10.2
Jersey City, NJ	14.4	2.2	0.09	8.6
Casper, WY	9.1	2.8	0.09	5.5
Freehold, NJ	38.3	16.0	0.44	35.3

Appendix IV. General Recommendations for Air Quality Improvement

Urban vegetation can directly and indirectly affect local and regional air quality by altering the urban atmosphere environment. Four main ways that urban trees affect air quality are (Nowak 1995):

- Temperature reduction and other microclimate effects
- Removal of air pollutants
- Emission of volatile organic compounds (VOC) and tree maintenance emissions
- Energy effects on buildings

The cumulative and interactive effects of trees on climate, pollution removal, and VOC and power plant emissions determine the impact of trees on air pollution. Cumulative studies involving urban tree impacts on ozone have revealed that increased urban canopy cover, particularly with low VOC emitting species, leads to reduced ozone concentrations in cities (Nowak 2000). Local urban management decisions also can help improve air quality.

Urban forest management strategies to help improve air quality include (Nowak 2000):

Strategy	Result
· · · · · · · · · · · · · · · · · · ·	
Increase the number of healthy trees	Increase pollution removal
Sustain existing tree cover	Maintain pollution removal levels
Maximize use of low VOC-emitting trees	Reduces ozone and carbon monoxide formation
Sustain large, healthy trees	Large trees have greatest per-tree effects
Use long-lived trees	Reduce long-term pollutant emissions from
	planting and removal
Use low maintenance trees	Reduce pollutants emissions from maintenance
	activities
Reduce fossil fuel use in maintaining vegetation	Reduce pollutant emissions
Plant trees in energy conserving locations	Reduce pollutant emissions from power plants
Plant trees to shade parked cars	Reduce vehicular VOC emissions
Supply ample water to vegetation	Enhance pollution removal and temperature
	reduction
Plant trees in polluted or heavily populated areas	Maximizes tree air quality benefits
Avoid pollutant-sensitive species	Improve tree health
Utilize evergreen trees for particulate matter	Year-round removal of particles

Section 9, ItemA.

Appendix V. Invasive Species of the Urban Forest

The following inventoried tree species were listed as invasive on the Washington invasive species list (Oregon Invasive Species Council 2014):

Species Name ^a	Number of Trees	% of Trees	Leaf Area	Percent Leaf Area
			(ac)	
Total	0	0.00	0.00	0.00

^aSpecies are determined to be invasive if they are listed on the state's invasive species list


Appendix VI. Potential Risk of Pests

Fifty-three insects and diseases were analyzed to quantify their potential impact on the urban forest. As each insect/disease is likely to attack different host tree species, the implications for {0} will vary. The number of trees at risk reflects only the known host species that are likely to experience mortality.

Code	Scientific Name	Common Name	Trees at Risk	Value
			(#)	(\$ millions)
AL	Phyllocnistis populiella	Aspen Leafminer	4,107	2.35
ALB	Anoplophora glabripennis	Asian Longhorned Beetle	66,215	125.94
ARCA	Neodothiora populina	Aspen Running Canker	0	0.00
ARD	Armillaria spp.	Armillaria Root Disease	75,703	266.33
BBD	Neonectria faginata	Beech Bark Disease	0	0.00
ВС	Sirococcus clavigignenti juglandacearum	Butternut Canker	105	0.02
BLD	Litylenchus crenatae mccannii	Beech Leaf Disease	0	0.00
BM	Euproctis chrysorrhoea	Browntail Moth	6,864	3.78
BOB	Tubakia iowensis	Bur Oak Blight	0	0.00
BSRD	Leptographium wageneri	Black Stain Root Disease	60,157	203.03
BWA	Adelges piceae	Balsam Woolly Adelgid	210	1.26
СВ	Cryphonectria parasitica	Chestnut Blight	0	0.00
DA	Discula destructiva	Dogwood Anthracnose	7,127	2.27
DBSR	Leptographium wageneri var. pseudotsugae	Douglas-fir Black Stain Root Disease	60,157	203.03
DED	Ophiostoma novo-ulmi	Dutch Elm Disease	0	0.00
DFB	Dendroctonus pseudotsugae	Douglas-Fir Beetle	48,440	185.81
EAB	Agrilus planipennis	Emerald Ash Borer	105	0.23
FE	Scolytus ventralis	Fir Engraver	48,650	185.96
FR	Cronartium quercuum f. sp. Fusiforme	Fusiform Rust	0	0.00
FTC	Malacosoma disstria	Forest Tent Caterpillar	10,491	21.72
GM	Lymantria dispar	Gypsy Moth	14,913	35.40
GSOB	Agrilus auroguttatus	Goldspotted Oak Borer	0	0.00
HRD	Heterobasidion irregulare/ occidentale	Heterobasidion Root Disease	87,420	284.66
HS	Neodiprion tsugae	Hemlock Sawfly	9,484	12.18
HWA	Adelges tsugae	Hemlock Woolly Adelgid	0	0.00
JPB	Dendroctonus jeffreyi	Jeffrey Pine Beetle	0	0.00
JPBW	Choristoneura pinus	Jack Pine Budworm	4,847	1.44
LAT	Choristoneura conflictana	Large Aspen Tortrix	13,005	24.15
LWD	Raffaelea lauricola	Laurel Wilt	1,813	0.55
MOB	Xyleborus monographus	Mediterranean Oak Borer	629	0.38
MPB	Dendroctonus ponderosae	Mountain Pine Beetle	3,454	8.95
NSE	lps perturbatus	Northern Spruce Engraver	1,011	2.68
OW	Ceratocystis fagacearum	Oak Wilt	0	0.00
PBSR	Leptographium wageneri var. ponderosum	Pine Black Stain Root Disease	2,023	1.30
POCRD	Phytophthora lateralis	Port-Orford-Cedar Root Disease	18,565	8.64

<u> </u>	Catanatific Name	Communication Name	Torres et D'ala	Section 9, ItemA.
Code	Scientific Name	Common Name	Trees at Risk	
			(#)	(\$ millions)
PSB	Tomicus piniperda	Pine Shoot Beetle	54,613	194.90
PSHB	Euwallacea nov. sp.	Polyphagous Shot Hole Borer	54,461	134.92
RPS	Matsucoccus resinosae	Red Pine Scale	0	0.00
SB	Dendroctonus rufipennis	Spruce Beetle	2,233	5.58
SBW	Choristoneura fumiferana	Spruce Budworm	49,661	188.71
SFM	subalpine fir mortality summary	Subalpine Fir Mortality	105	0.05
SLF	Lycorma delicatula	Spotted Lanternfly	22,400	12.08
SOD	Phytophthora ramorum	Sudden Oak Death	87,875	298.44
SPB	Dendroctonus frontalis	Southern Pine Beetle	16,773	22.95
SW	Sirex noctilio	Sirex Wood Wasp	6,173	9.09
TCD	Geosmithia morbida	Thousand Canker Disease	105	0.02
WBB	Dryocoetes confusus	Western Bark Beetle	0	0.00
WBBU	Acleris gloverana	Western Blackheaded Budworm	56,913	195.30
WFNPM	western five-needle pine	Western Five-Needle Pine	1,221	6.43
	mortality summary	Mortality		
WM	Operophtera brumata	Winter Moth	61,966	132.94
WPB	Dendroctonus brevicomis	Western Pine Beetle	0	0.00
WPBR	Cronartium ribicola	White Pine Blister Rust	1,221	6.43
WSB	Choristoneura occidentalis	Western Spruce Budworm	59,565	204.70

In the following graph, the pests are color coded according to the county's proximity to the pest occurrence in the United States. Red indicates that the pest is within the county; orange indicates that the pest is within 250 miles of the county; yellow indicates that the pest is within 750 miles of the county; and green indicates that the pest is outside of these ranges.



Note: points - Number of trees, bars - Replacement value

Based on the host tree species for each pest and the current range of the pest (Forest Health Technology Enterprise Team 2014), it is possible to determine what the risk is that each tree species in the urban forest could be attacked by an insect or disease.

			. Of disease.	_	_	1			_		_	_	_		_		_	_		_		_		_					_		_				_	_
Snn Rich	17:0	Kisk Weight	Species Name	٩٢	ALB	ARCA	ARD	BBD	BC	BLD	BM	BOB	BSRD	BWA	8	DA	DBSR	DED	DFB	EAB	田	FR	FTC	В	GSOB	HRD	£	HWA	JPB	JPBW	LAT	LWD	MOB	MPB	NSE	⊗
		32	Douglas fir																														П			
		29	Western white pine																														П		П	٦
		24	Subalpine fir																																	
		24	Norway spruce																														П		П	٦
		21	Lodgepole pine																																	
		19	Western hemlock																																	
		19	Mountain hemlock																																	
		16	Willow spp																																	
		12	Plum spp																																	
		12	Scots pine																																	
		12	Black cottonwood																																	
			Paper birch																																	
			Sitka spruce																																	
		10	Red alder																																	
		10	Blue spruce																																	
		10	European white birch																																	
		10	River birch																																	
		8	Western red cedar																														П			
		6	Apple spp																														П	П		
Ī	T	5	Bigleaf maple																														П			
		5	Northern white cedar																														П	П		٦
		5	Black poplar																														П	П		
		5	Black walnut																														П			
		4	Hinoki cypress																														П	П	П	
		4	Swiss mountain pine																														П	П		
Г	T	4	California laurel																														П	П	П	٦
		4	Port orford cedar																														П	П		
		4	Callery pear																														П	П	П	٦
			Pacific yew																																	
		4	Sweet cherry																																╗	\Box
		4	European mountain																														П	П		
			ash	L					L			L			L																		\bigsqcup			
		4	Oregon ash																																	
			Sweetgum																																	
		3	Japanese maple																																	
	Ī	3	Pacific dogwood																																	
		3	Red maple																																	
		3	Trident maple																																	
		2	Chinese parasoltree																																	
		2	Japanese flowering																																	
			cherry																														Ш			

																														ſ						
isk	しま	8	Γ	L	 	\Box	\Box			_1	<u>~</u>	۵	4			~		_	_					В			4	<u> </u>	>	\Box	S	Sect	ion	9, 1	lten	ıΑ.
Spp. Risk	Risk Weigh	Species Name	AL	ALB	ARC	ARD	8	BC	BLD	₩	BOB	BSRI	BW/	B	DA	DBSR	DED	DFB	EAB	뿐	표	FEC	В	eso	HR	HS	HWA	188	JPBV	\₹	ΓM	MO	ΔE	NS	ò	
		Coast redwood			П																												П			
	1	Bitter cherry			П																															
	1	Vine maple			П																															
	1	Camellia																															П			
	1	Oneseed hawthorn			П																															
	1	Kousa dogwood																															П			
	1	Flowering dogwood			П																												П			
	1	Japanese angelica tree	Г		П																												П			
	1	Atlas cedar			П																															
	1	Katsura tree	Г		П																												П			
	1	Babylon weeping																															П			
		willow																																		
	1	Southern magnolia																																		
	1	Black locust																																		
	1	Honeylocust																																		
	1	Common plum			I^{T}																															i

	um																				丄	丄
Spp. Risk	Risk Weight	Species Name	PBSR	POCRD	PSB	PSHB	RPS	SB	SBW	SFM	SLF	SOD	SPB	MS	ΩOI	WBB	WBBU	WFNPM	ΜM	WPB	WPBR	WSB
	32	Douglas fir																				
	29	Western white pine																				
	24	Subalpine fir																				
	24	Norway spruce																				
	21	Lodgepole pine																				
	19	Western hemlock																				
	19	Mountain hemlock																				
	16	Willow spp																				
	12	Plum spp																				
	12	Scots pine																				
	12	Black cottonwood																				П
	11	Paper birch																				
	11	Sitka spruce																				
	10	Red alder																				
	10	Blue spruce																				
	10	European white birch																				
	10	River birch																				
	8	Western red cedar																				
	6	Apple spp																				П
	5	Bigleaf maple																				
	5	Northern white cedar																				П
	5	Black poplar																				П
	5	Black walnut																				
	4	Hinoki cypress																				
	4	Swiss mountain pine																				
	4	California laurel																				П
	4	Port orford cedar																				
		i ort orrord cedar			_		_	_	_			<u> </u>	_				_	<u> </u>		<u> </u>	L	_

Section 9, ItemA.

. Risk	Risk Weight	rcies The	PBSR	CRD	SB	뫒	S _P	SB	BW	Ψ	iii	OO.	,PB	ΝS	<u> </u>	/BB	BBU	WFNPM	Σ×	VPB	PBR	WSB
Spp	≆ ≝	Spe	۵	Ы	Ĺ	ثق	Ľ		s	S	Ľ.	S	<u></u>	Ľ		^	≥	W	۱	>	≥	_
		Callery pear																				Ш
		Pacific yew																			L	
		Sweet cherry																			L	
		European mountain																				
		ash																				Ш
		Oregon ash														L					L	Ш
		Sweetgum																				
		Japanese maple																			L	
		Pacific dogwood																				
		Red maple																				
	3	Trident maple																				
	2	Chinese parasoltree																				
	2	Japanese flowering																				
		cherry																			L	
	2	Coast redwood																				
	1	Bitter cherry																				
	1	Vine maple																				
	1	Camellia																				
	1	Oneseed hawthorn																				
	1	Kousa dogwood																				
	1	Flowering dogwood																				
	1	Japanese angelica tree																				
	1	Atlas cedar																				
	1	Katsura tree																				
	1	Babylon weeping																				
		willow		L				L														
	1	Southern magnolia																				
	1	Black locust																				
	1	Honeylocust																				
	1	Common plum																				

Note:

Species that are not listed in the matrix are not known to be hosts to any of the pests analyzed.

Species Risk:

- Red indicates that tree species is at risk to at least one pest within county
- Orange indicates that tree species has no risk to pests in county, but has a risk to at least one pest within 250 miles from the county
- Yellow indicates that tree species has no risk to pests within 250 miles of county, but has a risk to at least one pest that is 250 and 750 miles from the county
- Green indicates that tree species has no risk to pests within 750 miles of county, but has a risk to at least one pest that is greater than 750 miles from the county

Risk Weight:

Numerical scoring system based on sum of points assigned to pest risks for species. Each pest that could attack tree species is scored as 4 points if red, 3 points if orange, 2 points if yellow and 1 point if green.

Pest Color Codes:

- Red indicates pest is within King county
- Red indicates pest is within 250 miles county
- Yellow indicates pest is within 750 miles of King county
- Green indicates pest is outside of these ranges

Additional References for i-Tree Model, Analysis, and Appendices

Abdollahi, K.K.; Ning, Z.H.; Appeaning, A., eds. 2000. Global climate change and the urban forest. Baton Rouge, LA: GCRCC and Franklin Press. 77 p.

Baldocchi, D. 1988. A multi-layer model for estimating sulfur dioxide deposition to a deciduous oak forest canopy. Atmospheric Environment. 22: 869-884.

Baldocchi, D.D.; Hicks, B.B.; Camara, P. 1987. A canopy stomatal resistance model for gaseous deposition to vegetated surfaces. Atmospheric Environment. 21: 91-101.

Bidwell, R.G.S.; Fraser, D.E. 1972. Carbon monoxide uptake and metabolism by leaves. Canadian Journal of Botany. 50: 1435-1439.

British Columbia Ministry of Water, Land, and Air Protection. 2005. Residential wood burning emissions in British Columbia. British Columbia.

Broecker, W.S. 1970. Man's oxygen reserve. Science 168(3939): 1537-1538.

Bureau of Transportation Statistics. 2010. Estimated National Average Vehicle Emissions Rates per Vehicle by Vehicle Type using Gasoline and Diesel. Washington, DC: Burea of Transportation Statistics, U.S. Department of Transportation. Table 4-43.

California Air Resources Board. 2013. Methods to Find the Cost-Effectiveness of Funding Air Quality Projects. Table 3 Average Auto Emission Factors. CA: California Environmental Protection Agency, Air Resources Board.

Carbon Dioxide Information Analysis Center. 2010. CO2 Emissions (metric tons per capita). Washington, DC: The World Bank.

Cardelino, C.A.; Chameides, W.L. 1990. Natural hydrocarbons, urbanization, and urban ozone. Journal of Geophysical Research. 95(D9): 13,971-13,979.

Ciesla, W. M.; Kruse, J. J. 2009. Large Aspen Tortrix. Forest Insect & Disease Leaflet 139. Washington, DC: U. S. Department of Agriculture, Forest Service. 8 p.

DeMars, C. J., Jr.; Roettgering, B. H. 1982. Western Pine Beetle. Forest Insect & Disease Leaflet 1. Washington, DC: U.S. Department of Agriculture, Forest Service. 8 p.

Eastern Forest Environmental Threat Assessment Center. Dutch Elm Disease. http://threatsummary.forestthreats.org/threats/threatSummaryViewer.cfm?threatID=43

Energy Information Administration. 1994. Energy Use and Carbon Emissions: Non-OECD Countries. Washington, DC: Energy Information Administration, U.S. Department of Energy.

Energy Information Administration. 2013. CE2.1 Fuel consumption totals and averages, U.S. homes. Washington, DC: Energy Information Administration, U.S. Department of Energy.

Energy Information Administration. 2014. CE5.2 Household wood consumption. Washington, DC: Energy Information Administration, U.S. Department of Energy.

Federal Highway Administration. 2013. Highway Statistics 2011. Washington, DC: Federal Highway Administration, U.S. Department of Transportation. Table VM-1.

Fellin, D. G.; Dewey, J. E. 1986. Western Spruce Budworm. Forest Insect & Disease Leaflet 53. Washington, DC: U.S. Department of Agriculture, Forest Service. 10 p.

Ferrell, G. T. 1986. Fir Engraver. Forest Insect & Disease Leaflet 13. Washington, DC: U. S. Department of Agriculture, Forest Service. 8 p.

Forest Health Technology Enterprise Team. 2014. 2012 National Insect & Disease Risk Maps/Data. Fort Collins, CO: U.S. Department of Agriculture, Forest Service. http://www.fs.fed.us/foresthealth/technology/nidrm2012.shtml

Georgia Forestry Commission. 2009. Biomass Energy Conversion for Electricity and Pellets Worksheet. Dry Branch, GA: Georgia Forestry Commission.

Gibson, K.; Kegley, S.; Bentz, B. 2009. Mountain Pine Beetle. Forest Insect & Disease Leaflet 2. Washington, DC: U. S. Department of Agriculture, Forest Service. 12 p.

Heirigs, P.L.; Delaney, S.S.; Dulla, R.G. 2004. Evaluation of MOBILE Models: MOBILE6.1 (PM), MOBILE6.2 (Toxics), and MOBILE6/CNG. Sacramento, CA: National Cooperative Highway Research Program, Transportation Research Board.

Hirabayashi, S. 2011. Urban Forest Effects-Dry Deposition (UFORE-D) Model Enhancements, http://www.itreetools.org/eco/resources/UFORE-D enhancements.pdf

Hirabayashi, S. 2012. i-Tree Eco Precipitation Interception Model Descriptions, http://www.itreetools.org/eco/resources/iTree Eco Precipitation Interception Model Descriptions V1 2.pdf

Hirabayashi, S.; Kroll, C.; Nowak, D. 2011. Component-based development and sensitivity analyses of an air pollutant dry deposition model. Environmental Modeling and Software. 26(6): 804-816.

Hirabayashi, S.; Kroll, C.; Nowak, D. 2012. i-Tree Eco Dry Deposition Model Descriptions V 1.0

Holsten, E.H.; Thier, R.W.; Munson, A.S.; Gibson, K.E. 1999. The Spruce Beetle. Forest Insect & Disease Leaflet 127. Washington, DC: U.S. Department of Agriculture, Forest Service. 12 p.

Interagency Working Group on Social Cost of Carbon, United States Government. 2015. Technical Support Document: Technical Update of the Social Cost of Carbon for Regulatory Impact Analysis Under Executive Order 12866. http://www.whitehouse.gov/sites/default/files/omb/inforeg/scc-tsd-final-july-2015.pdf

Layton, M. 2004. 2005 Electricity Environmental Performance Report: Electricity Generation and Air Emissions. CA: California Energy Commission.

Section 9. ItemA.

Leonardo Academy. 2011. Leonardo Academy's Guide to Calculating Emissions Including Emission Factor Prices. Madison, WI: Leonardo Academy Inc.

Liebhold, A. 2010 draft. Personal communication on the geographic distribution of forest pest species.

Lovett, G.M. 1994. Atmospheric deposition of nutrients and pollutants in North America: an ecological perspective. Ecological Applications. 4: 629-650.

McPherson, E.G.; Maco, S.E.; Simpson, J.R.; Peper, P.J.; Xiao, Q.; VanDerZanden, A.M.; Bell, N. 2002. Western Washington and Oregon Community Tree Guide: Benefits, Costs, and Strategic Planting. International Society of Arboriculture, Pacific Northwest, Silverton, OR.

McPherson, E.G.; Simpson, J.R. 1999. Carbon dioxide reduction through urban forestry: guidelines for professional and volunteer tree planters. Gen. Tech. Rep. PSW-171. Albany, CA: U.S. Department of Agriculture, Forest Service, Pacific Southwest Research Station. 237 p.

McPherson, E.G.; Simpson, J.R.; Peper, P.J.; Crowell, A.M.N.; Xiao, Q. 2010. Northern California coast community tree guide: benefits, costs, and strategic planting. PSW-GTR-228. Gen. Tech. Rep. PSW-GTR-228. U.S. Department of Agriculture, Forest Service, Pacific Southwest Research Station, Albany, CA.

McPherson, E.G.; Simpson, J.R.; Peper, P.J.; Gardner, S.L.; Vargas, K.E.; Maco, S.E.; Xiao, Q. 2006a. Coastal Plain Community Tree Guide: Benefits, Costs, and Strategic Planting PSW-GTR-201. USDA Forest Service, Pacific Southwest Research Station, Albany, CA.

McPherson, E.G.; Simpson, J.R.; Peper, P.J.; Gardner, S.L.; Vargas, K.E.; Xiao, Q. 2007. Northeast community tree guide: benefits, costs, and strategic planting.

McPherson, E.G.; Simpson, J.R.; Peper, P.J.; Maco, S.E.; Gardner, S.L.; Cozad, S.K.; Xiao, Q. 2006b. Midwest Community Tree Guide: Benefits, Costs and Strategic Planting PSW-GTR-199. U.S. Department of Agriculture, Forest Service, Pacific Southwest Research Station, Albany, CA.

McPherson, E.G.; Simpson, J.R.; Peper, P.J.; Maco, S.E.; Gardner, S.L.; Vargas, K.E.; Xiao, Q. 2006c. Piedmont Community Tree Guide: Benefits, Costs, and Strategic Planting PSW-GTR 200. U.S. Department of Agriculture, Forest Service, Pacific Southwest Research Station, Albany, CA.

McPherson, E.G.; Simpson, J.R.; Peper, P.J.; Maco, S.E.; Xiao Q.; Mulrean, E. 2004. Desert Southwest Community Tree Guide: Benefits, Costs and Strategic Planting. Phoenix, AZ: Arizona Community Tree Council, Inc. 81:81.

McPherson, E.G.; Simpson, J.R.; Peper, P.J.; Scott, K.I.; Xiao, Q. 2000. Tree Guidelines for Coastal Southern California Communities. Local Government Commission, Sacramento, CA.

McPherson, E.G.; Simpson, J.R.; Peper, P.J.; Xiao, Q. 1999. Tree Guidelines for San Joaquin Valley Communities. Local Government Commission, Sacramento, CA.

McPherson, E.G.; Simpson, J.R.; Peper, P.J.; Xiao, Q.; Maco, S.E.; Hoefer, P.J. 2003. Northern Mountain and Prairie Community Tree Guide: Benefits, Costs and Strategic Planting. Center for Urban Forest Research, USDA Forest Service, Pacific Southwest Research Station, Albany, CA.

McPherson, E.G.; Simpson, J.R.; Peper, P.J.; Xiao, Q.; Pittenger, D.R.; Hodel, D.R. 2001. Tree Guidelines for Inland Empire Communities. Local Government Commission, Sacramento, CA.

Section 9. ItemA.

Murray, F.J.; Marsh L.; Bradford, P.A. 1994. New York State Energy Plan, vol. II: issue reports. Albany, State Energy Office.

National Invasive Species Information Center. 2011. Beltsville, MD: U.S. Department of Agriculture, National Invasive Species Information Center. http://www.invasivespeciesinfo.gov/plants/main.shtml

Nicholls, T. H.; Anderson, R. L. 1977. How to Identify White Pine Blister Rust and Remove Cankers. St. Paul, MN: U.S. Department of Agriculture, Forest Service, Northeastern Area State and Private Forestry

Nowak, D.J. 1994. Atmospheric carbon dioxide reduction by Chicago's urban forest. In: McPherson, E.G.; Nowak, D.J.; Rowntree, R.A., eds. Chicago's urban forest ecosystem: results of the Chicago Urban Forest Climate Project. Gen. Tech. Rep. NE-186. Radnor, PA: U.S. Department of Agriculture, Forest Service, Northeastern Forest Experiment Station: 83-94.

Nowak, D.J. 1995. Trees pollute? A "TREE" explains it all. In: Proceedings of the 7th National Urban Forestry Conference. Washington, DC: American Forests: 28-30.

Nowak, D.J. 2000. The interactions between urban forests and global climate change. In: Abdollahi, K.K.; Ning, Z.H.; Appeaning, A., eds. Global Climate Change and the Urban Forest. Baton Rouge, LA: GCRCC and Franklin Press: 31-44.

Nowak, D.J., Hirabayashi, S., Bodine, A., Greenfield, E. 2014. Tree and forest effects on air quality and human health in the United States. Environmental Pollution. 193:119-129.

Nowak, D.J., Hirabayashi, S., Bodine, A., Hoehn, R. 2013. Modeled PM2.5 removal by trees in ten U.S. cities and associated health effects. Environmental Pollution. 178: 395-402.

Nowak, D.J.; Civerolo, K.L.; Rao, S.T.; Sistla, S.; Luley, C.J.; Crane, D.E. 2000. A modeling study of the impact of urban trees on ozone. Atmospheric Environment. 34: 1601-1613.

Nowak, D.J.; Crane, D.E. 2000. The Urban Forest Effects (UFORE) Model: quantifying urban forest structure and functions. In: Hansen, M.; Burk, T., eds. Integrated tools for natural resources inventories in the 21st century. Proceedings of IUFRO conference. Gen. Tech. Rep. NC-212. St. Paul, MN: U.S. Department of Agriculture, Forest Service, North Central Research Station: 714-720.

Nowak, D.J.; Crane, D.E.; Dwyer, J.F. 2002a. Compensatory value of urban trees in the United States. Journal of Arboriculture. 28(4): 194 - 199.

Nowak, D.J.; Crane, D.E.; Stevens, J.C.; Hoehn, R.E. 2005. The urban forest effects (UFORE) model: field data collection manual. V1b. Newtown Square, PA: U.S. Department of Agriculture, Forest Service, Northeastern Research Station, 34 p. http://www.fs.fed.us/ne/syracuse/Tools/downloads/UFORE_Manual.pdf

Nowak, D.J.; Crane, D.E.; Stevens, J.C.; Ibarra, M. 2002b. Brooklyn's urban forest. Gen. Tech. Rep. NE-290. Newtown Square, PA: U.S. Department of Agriculture, Forest Service, Northeastern Research Station. 107 p.

Nowak, D.J.; Dwyer, J.F. 2000. Understanding the benefits and costs of urban forest ecosystems. In: Kuser, John, ed. Handbook of urban and community forestry in the northeast. New York, NY: Kluwer Academics/Plenum: 11-22.

Nowak, D.J.; Hoehn, R.; Crane, D. 2007. Oxygen production by urban trees in the United States. Arboriculture & Urban Forestry 33(3):220-226.

Nowak, D.J.; Hoehn, R.E.; Crane, D.E.; Stevens, J.C.; Walton, J.T; Bond, J. 2008. A ground-based method of assessing urban forest structure and ecosystem services. Arboriculture and Urban Forestry. 34(6): 347-358.

Nowak, D.J.; Stevens, J.C.; Sisinni, S.M.; Luley, C.J. 2002c. Effects of urban tree management and species selection on atmospheric carbon dioxide. Journal of Arboriculture. 28(3): 113-122.

Oregon Invasive Species Council. 2014. 100 Worst List. OR: Oregon Invasive Species Council. http://www.oregoninvasivespeciescouncil.org/media/100_worst.pdf>

Peper, P.J.; McPherson, E.G.; Simpson, J.R.; Albers, S.N.; Xiao, Q. 2010. Central Florida community tree guide: benefits, costs, and strategic planting. Gen. Tech. Rep. PSW-GTR-230. U.S. Department of Agriculture, Forest Service, Pacific Southwest Research Station, Albany, CA.

Peper, P.J.; McPherson, E.G.; Simpson, J.R.; Vargas, K.E.; Xiao Q. 2009. Lower Midwest community tree guide: benefits, costs, and strategic planting. PSW-GTR-219. Gen. Tech. Rep. PSW-GTR-219. U.S. Department of Agriculture, Forest Service, Pacific Southwest Research Station, Albany, CA.

Schmitz, R. F.; Gibson, K. E. 1996. Douglas-fir Beetle. Forest Insect & Disease Leaflet 5. R1-96-87. Washington, DC: U. S. Department of Agriculture, Forest Service. 8 p.

U.S. Environmental Protection Agency. 2010. Light-Duty Vehicle Greenhouse Gas Emission Standards and Corporate Average Fuel Economy Standards. Washington, DC: U.S. Environmental Protection Agency. EPA-420-R-10-012a

U.S. Environmental Protection Agency. 2015. The social cost of carbon. http://www.epa.gov/climatechange/EPAactivities/economics/scc.html

van Essen, H.; Schroten, A.; Otten, M.; Sutter, D.; Schreyer, C.; Zandonella, R.; Maibach, M.; Doll, C. 2011. External Costs of Transport in Europe. Netherlands: CE Delft. 161 p.

Vargas, K.E.; McPherson, E.G.; Simpson, J.R.; Peper, P.J.; Gardner, S.L.; Xiao, Q. 2007a. Interior West Tree Guide.

Vargas, K.E.; McPherson, E.G.; Simpson, J.R.; Peper, P.J.; Gardner, S.L.; Xiao, Q. 2007b. Temperate Interior West Community Tree Guide: Benefits, Costs, and Strategic Planting.

Vargas, K.E.; McPherson, E.G.; Simpson, J.R.; Peper, P.J.; Gardner, S.L.; Xiao, Q. 2008. Tropical community tree guide: benefits, costs, and strategic planting. PSW-GTR-216. Gen. Tech. Rep. PSW-GTR-216. U.S. Department of Agriculture, Forest Service, Pacific Southwest Research Station, Albany, CA.

Worrall, J.J. 2007. Chestnut Blight. Forest and Shade Tree Pathology. http://www.forestpathology.org/dis_chestnut.html

Zinke, P.J. 1967. Forest interception studies in the United States. In: Sopper, W.E.; Lull, H.W., eds. Forest Hydrology. Oxford, UK: Pergamon Press: 137-161.

Project Name	Address	Review Date	Permit Purpose	Project Status
Patton	18711 35th Ave NE	10/13/2023	Removal to mitigate risk	Responses received
Fobes	19750 41st Ave NE	10/13/2023	Removal to mitigate risk	minor update requested
Headwaters	18948 Forest Park Drive	10/13/2023	Right-of-way tree removal	Approved
Seely	16730 Shore Drive NE	11/2/2023	Nuisance tree removal	waiting for response
Lakeview Townhomes	3803 NE 155th St	11/3/2023	Major develoment activity	waiting for response
Skalenakis	4728 NE 204th St	11/6/2023	Removal to mitigate risk	Needs replanting plan
Oakes	Parcel 4017110480	11/6/2023	Removal to mitigate risk	Approved
McLeod	3781 NE 185th St	11/6/2023	Removal to mitigate risk	Needs replanting plan
Caba	33rd Ave NE	11/7/2023	Major develoment activity	waiting for response
Brito	18325 28th Place	11/17/2023	Minor development activity	Director considering variance
Crandall	19946 47th Ave NE	11/17/2023	Removal to mitigate risk	Approved
Rollinger	4510 NE 190th Ct	11/17/2023	Removal to mitigate risk	in conflict with ECA rules
Larkin	5323 NE 180th Street	11/17/2023	Minor development activity	waiting for response